

Container Label

GROUP	3	11	FUNGICIDE
-------	---	----	-----------

**HEADLINE® AMP**  
**Fungicide**

SOLUTION

COMMERCIAL - AGRICULTURAL

Broad spectrum fungicide for use in cereals and corn

**ACTIVE INGREDIENTS:** Pyraclostrobin.....146 g/L  
Metconazole ..... 55 g/L

Contains 2-methyl-4-isothiazolin-3-one and 1,2-benzisothiazolin-3-one, both at 0.0025%, as preservatives

**OR**

Contains 2-methyl-4-isothiazolin-3-one and 1,2-benzisothiazolin-3-one, both at 0.0075% and 2-bromo-2-nitropropane-1,3-diol at 0.024%, as preservatives

**OR**

Contains 1,2-benzisothiazolin-3-one at 0.0400%, 5-chloro-2-methyl-4-isothiazolin-3-one at 0.0011%, 2-methyl-4-isothiazolin-3-one at 0.0004%, and 2-bromo-2-nitropropane-1,3-diol at 0.0240%, as preservatives

**REGISTRATION NO. 32189**

**PEST CONTROL PRODUCTS ACT**



**WARNING**

**POISON**

**SKIN IRRITANT**

**READ THE LABEL AND ATTACHED BROCHURE BEFORE USING**  
**KEEP OUT OF REACH OF CHILDREN**

**IN CASE OF EMERGENCY ENDANGERING LIFE OR PROPERTY**  
**INVOLVING THIS PRODUCT, CALL DAY OR NIGHT,**  
**1-800-454-2673**

**NET CONTENTS:** 0.100 L to bulk

BASF Agricultural Solutions Canada Inc.  
510 - 28 Quarry Park Blvd. SE  
Calgary, Alberta T2C 5P9  
1-877-371-2273

HEADLINE is a registered trade-mark of BASF Corporation, used under license by BASF Agricultural Solutions Canada Inc.

## PRECAUTIONS

1. **KEEP OUT OF REACH OF CHILDREN.**
2. May be fatal or poisonous if swallowed. Do not take internally. Harmful if inhaled. Avoid inhaling/breathing dusts, sprays, etc.
3. Causes moderate eye and skin irritation. DO NOT get in eyes or on skin.
4. Wash thoroughly after handling and before eating, drinking or smoking.
5. **All mixer/loaders and applicators (except aerial applicators):** Wear coveralls over long sleeved shirt, long pants, chemical resistant gloves, socks and footwear during mixing/loading, application, clean-up and repair. In addition, wear goggles/face shield during mixing/loading, clean-up and repair. Gloves are not required during application.

**Aerial applicators:** Wear long sleeved shirt, long pants, socks and footwear during application.

Use closed cab tractor for ground boom application. A closed cab is a chemical resistant barrier that completely surrounds the occupant of the cab and prevents contact with the pesticide or treated surfaces outside the cab.

6. If clothing becomes contaminated, remove and wash separately from household laundry before reuse.
7. Clean spray equipment thoroughly after use.
8. Apply only to agricultural crops when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

## FIRST AID

**If swallowed:** Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

## **TOXICOLOGICAL INFORMATION**

The patient should be treated symptomatically.

## **ENVIRONMENTAL PRECAUTIONS**

Toxic to aquatic organisms, non-target terrestrial plants and small wild mammals.

Observe buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g. sandy soil) and/or the depth to the water table is shallow.

## **STORAGE**

1. Store in original tightly closed container. Protect from freezing.
2. Do not ship or store near food, feed, seed and fertilizers.
3. Store in cool, dry, locked, well-ventilated area without floor drain.

## **DISPOSAL**

### **For Recyclable Containers**

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for any further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

**For Non-Returnable Containers**

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

**For Returnable-Refillable Containers**

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

**NOTICE TO USER**

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

Booklet

GROUP	3	11	FUNGICIDE
-------	---	----	-----------

**HEADLINE® AMP  
Fungicide**

SOLUTION

COMMERCIAL - AGRICULTURAL

Broad spectrum fungicide for use in cereals and corn

**ACTIVE INGREDIENTS:** Pyraclostrobin.....146 g/L  
Metconazole ..... 55 g/L

Contains 2-methyl-4-isothiazolin-3-one and 1,2-benzisothiazolin-3-one, both at 0.0025%, as preservatives

**OR**

Contains 2-methyl-4-isothiazolin-3-one and 1,2-benzisothiazolin-3-one, both at 0.0075% and 2-bromo-2-nitropropane-1,3-diol at 0.024%, as preservatives

**OR**

Contains 1,2-benzisothiazolin-3-one at 0.0400%, 5-chloro-2-methyl-4-isothiazolin-3-one at 0.0011%, 2-methyl-4-isothiazolin-3-one at 0.0004%, and 2-bromo-2-nitropropane-1,3-diol at 0.0240%, as preservatives

**REGISTRATION NO. 32189**

**PEST CONTROL PRODUCTS ACT**



**WARNING**

**POISON**

**SKIN IRRITANT**

**READ THE LABEL AND ATTACHED BROCHURE BEFORE USING  
KEEP OUT OF REACH OF CHILDREN**

**IN CASE OF EMERGENCY ENDANGERING LIFE OR PROPERTY  
INVOLVING THIS PRODUCT, CALL DAY OR NIGHT,  
1-800-454-2673**

**NET CONTENTS:** 0.100 L to bulk

BASF Agricultural Solutions Canada Inc.  
510 - 28 Quarry Park Blvd. SE  
Calgary, Alberta T2C 5P9  
1-877-371-2273

HEADLINE is a registered trade-mark of BASF Corporation, used under license by BASF Agricultural Solutions Canada Inc.

## GENERAL INFORMATION

This package contains **HEADLINE AMP**, a 201 g/L suspension concentrate. The active ingredient in **HEADLINE AMP** belongs to the triazole and strobilurin class of fungicides.

## DIRECTIONS FOR USE (See specific sections for each crop group)

For control of the leaf diseases listed below, apply 0.5 to 1.0 L of **HEADLINE AMP** per hectare in cereals **OR** 0.75 to 1.0 L of **HEADLINE AMP** per hectare in corn. **HEADLINE AMP** can be applied with either ground or aerial application equipment. Equipment should be checked frequently for proper application.

## TANK MIXING

This product may be tank mixed with a fertilizer, a supplement or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact BASF Agricultural Solutions Canada Inc. ("BASF") at 1-877-371-2273 for information before mixing any pesticide that is not specifically recommended on this label.

Not all varieties and cultivars have been tested with possible tank mix combinations. Local conditions can also influence crop tolerance and may not match those under which BASF has conducted testing. Physical incompatibility, reduced disease control, or crop injury may result from mixing **HEADLINE AMP** with other products. Therefore, before using any tank mix (fungicides, insecticides, herbicides, liquid fertilizers, biological control products, adjuvants, and additives), test the combination on a small portion of the crop to be treated to ensure that a phytotoxic response will not occur as a result of application. Consult a BASF representative for more information concerning additives.

### Tank Mix Compatibility Determination

To determine compatibility, mix all components of the finished spray in proportionate quantities in a small jar. Add components to the test jar in the following order: wettable powders and water dispersible granules, liquid flowables and suspensions, emulsifiable concentrate formulations, and solutions. If the mixture does not ball-up or form flakes, sludge, jelly, oily films or layers, or other precipitates within 2 hours after mixing and can easily be resuspended with agitation, then the tested spray-mix is compatible. Always follow the most restrictive label regarding any precautions when tank mixing.

### Cereals Application Rate Table (Ground and aerial application)

Crop	Disease	Application Rate (L/ha)
Barley	Net blotch ( <i>Pyrenophora teres</i> ) Spot blotch ( <i>Cochliobolus sativus</i> ) Scald ( <i>Rhynchosporium secalis</i> ) Stripe rust ( <i>Puccinia striiformis</i> )	0.5 - 1.0
Wheat (all types) Triticale	Tan spot ( <i>Pyrenophora tritici-repentis</i> ) Septoria leaf spot ( <i>Septoria tritici</i> <b>OR</b> <i>Stagonospora nodorum</i> ) Leaf rust ( <i>Puccinia recondita</i> ) Spot blotch ( <i>Cochliobolus sativus</i> ) Stripe rust ( <i>Puccinia striiformis</i> ) Powdery mildew ( <i>Erysiphe graminis</i> f. <i>sp. tritici</i> )	
Oat	Crown rust ( <i>Puccinia coronata</i> )	
Rye	Leaf rust ( <i>Puccinia recondita</i> ) Powdery mildew ( <i>Erysiphe graminis</i> )	

#### Application Directions

##### Leaf Diseases in Cereals

For optimal leaf disease control, begin applications of **HEADLINE AMP** prior to disease development or at the onset of disease symptoms. Use the higher rate and shorter interval when disease pressure is high.

Minimum spray interval is 10 - 14 days.

##### Resistance Management

DO NOT make more than two (2) applications of **HEADLINE AMP** per season. Do not make sequential applications of **HEADLINE AMP**. If disease persists or weather conditions are favorable for disease development, make a second application 10-14 days later with a fungicide that contains a different mode of action.

### Corn Application Rate Table (Ground and aerial application)

Crop	Disease	Application Rate (L/ha)
Corn (field, pop, sweet, seed)	Anthracnose ( <i>Colletotrichum graminicola</i> ) Common rust ( <i>Puccinia sorghi</i> ) Gray leaf spot ( <i>Cercospora zeae-maydis</i> ) Northern leaf blight ( <i>Setosphaeria turcica</i> ) Eye spot ( <i>Aureobasidium zeae</i> )	0.75 - 1.0
<p><b>Application Directions</b></p> <p>For optimal disease control, begin applications of <b>HEADLINE AMP</b> prior to disease development. Use the higher rate and shorter interval when disease pressure is high.</p> <p>Do not make sequential applications of <b>HEADLINE AMP</b>. If disease persists or weather conditions are favourable for disease development make a second application 10 - 14 days later, with a fungicide that contains a different mode of action.</p> <p>DO NOT make more than two (2) applications of <b>HEADLINE AMP</b> per season.</p>		

### Corn Application Rate Table (Ground and aerial application) – Tank Mixture of 0.75 L/ha HEADLINE AMP plus 0.5 L/ha CARAMBA Fungicide for suppression of Fusarium and Gibberella ear rots and control of foliar diseases

Crop	Disease	Application Rate (L/ha)	
		HEADLINE AMP	CARAMBA Fungicide
Corn (field, pop, sweet, seed)	For suppression of fusarium and gibberella ear rots ( <i>Fusarium graminearum</i> and <i>Gibberella zeae</i> )	0.75	0.5
	For control of: Anthracnose ( <i>Colletotrichum graminicola</i> ) Common rust ( <i>Puccinia sorghi</i> ) Gray leaf spot ( <i>Cercospora zeae-maydis</i> ) Northern leaf blight ( <i>Setosphaeria turcica</i> ) Eye spot ( <i>Aureobasidium zeae</i> )		
<p><b>Application Directions</b></p> <p><b>Suppression of Fusarium and Gibberella ear rots in Corn</b></p> <p>The risk of Fusarium and Gibberella ear rots in corn is greater when the weather is warm and wet during silking. The application of <b>HEADLINE AMP</b> plus <b>CARAMBA Fungicide</b> (PCP Reg. No. 29767) for suppression of Fusarium and Gibberella ear rots should be considered when these environmental conditions are forecasted.</p> <p>To maximize yield and suppress Fusarium and Gibberella ear rots, apply <b>HEADLINE AMP</b> tank mixed with <b>CARAMBA Fungicide</b> when the crop is between silking (GS 63) and silk</p>			



Crop	Disease	Application Rate (L/ha)	
		HEADLINE AMP	CARAMBA Fungicide
<p>browning (GS 67). Application of <b>HEADLINE AMP</b> tank mixed with <b>CARAMBA Fungicide</b> will reduce the Fusarium toxin (DON) in harvested grain.</p> <p>DO NOT make more than one (1) application of <b>HEADLINE AMP + CARAMBA Fungicide</b> tank mix per season.</p> <p>Spray coverage: It is important to have good spray coverage on the silks to ensure optimum efficacy.</p> <p><b>Control of foliar leaf diseases in Corn</b></p> <p><b>HEADLINE AMP</b> tank mixed with <b>CARAMBA Fungicide</b> applied to suppress Fusarium and Gibberella ear rots will also control listed foliar leaf diseases. Optimal leaf disease control occurs when <b>HEADLINE AMP</b> tank mixed with <b>CARAMBA Fungicide</b> is applied prior to disease development or at the onset of disease symptoms.</p>			

### Field sprayer application

**DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural and Biological Engineers (ASABE) medium classification. Boom height must be 60 cm or less above the crop or ground.

### Aerial application

**DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply when wind speed is greater than 16 km/h at flying height at the site of application. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural and Biological Engineers (ASABE) medium classification. To reduce drift caused by turbulent wingtip vortices, the nozzle distribution along the spray boom length **MUST NOT** exceed 65% of the wing- or rotor-span.

### Spray Buffer zones

Use of the following spray methods or equipment **DOES NOT** require a spray buffer zone: hand-held or backpack sprayer and spot treatment.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of Application	Crop		Spray Buffer Zones (metres) Required for the Protection of:			
			Freshwater Habitat of Depths:		Estuarine/Marine Habitats of Depths:	
			Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m
Field sprayer	Barley, wheat (all types), triticale, oats, rye, corn		10	1	1	1
Aerial (Fixed and rotary wing)	Barley, wheat (all types), triticale, oats, rye, corn	For rates up to 0.75 L/ha	85	10	10	1
	Barley, wheat (all types), triticale, oats, rye, corn	For rates above 0.75 L/ha to 1.0 L/ha	125	15	10	3

When tank mixes are permitted, consult the labels of the tank mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASABE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticides portion of the Canada.ca website.

### General Restrictions

As this product is not registered for the control of pests in aquatic systems, **DO NOT** use to control aquatic pests.

**DO NOT** contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

### Ground Application

Apply **HEADLINE AMP** at rates listed in application rate and timing table (crop specific) when conditions are favourable for the development of disease. Use a minimum water volume of 100 L/ha. Ensure thorough coverage of foliage. Consult nozzle manufacturers recommendation for spray pressures for specific nozzles.

### Aerial Application

Apply **HEADLINE AMP** at rates listed in the application rate and timing tables (crop specific) when conditions are favourable for the development of disease. Use a minimum water volume of 50 L/ha. Ensure thorough coverage of foliage. Consult nozzle manufacturers recommendation for spray pressures for specific nozzles.

### Instructions for Aerial Applications

Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

### **Use Precautions**

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the National Aerial Pesticide Application Manual, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

Do not apply to any body of water. Avoid drifting of spray onto any body of water or other non-target areas. Specified buffer zones should be observed.

Coarse sprays are less likely to drift, therefore, avoid combinations of pressure and nozzle type that will result in fine particles (mist). Do not apply during periods of dead calm or when wind velocity and direction pose a risk of spray drift. Do not spray when the wind is blowing towards a nearby sensitive crop, garden, terrestrial habitat (such as shelter-belt) or aquatic habitat.

### **Operator Precautions**

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

### **Product Specific Precautions**

Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-877-371-2273 or obtain technical advice from the distributor or your provincial agricultural representative.

### **Buffer Zone for Aerial Application**

Aerial drift is increased under certain meteorological conditions. Do not apply during periods of dead calm, when winds are gusty or when wind speed is greater than 16 km/hour at flying height at the site of application.

For the protection of non-target habitats, overspray or drift to sensitive habitats must be avoided. A buffer zone of 5 metres is required between the downwind point of direct application and the closest edge of sensitive terrestrial habitats including grasslands, forested areas, shelter belts, woodlots, hedgerows, pastures, rangelands and shrublands. The buffer zones specified in the table below are required between the downwind point of direct application and the closest edge of sensitive aquatic habitats such as lakes, rivers, sloughs, ponds, coulees, prairie potholes, creeks, marshes, streams, reservoirs and wetlands. Do not contaminate these habitats when cleaning and rinsing spray equipment or containers.

## RESISTANCE MANAGEMENT RECOMMENDATIONS

### Mode of Action

Metconazole, one of the active ingredients of **HEADLINE AMP**, is classified by the Fungicide Resistance Action Committee (FRAC) as a Target Site of Action Group 3 Fungicide. Pyraclostrobin, the other active ingredient of **HEADLINE AMP**, is classified by FRAC as a Target Site of Action Group 11 Fungicide.

### Resistance management

**HEADLINE AMP** contains metconazole, a Group 3 fungicide and pyraclostrobin, a Group 11 fungicide. Any fungal population may contain individuals naturally resistant to Group 3 and 11 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to a site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance management strategies should be followed.

The following recommendations may be considered to delay the development of fungicide resistance:

1. **Applications:** Where possible, rotate the use of **HEADLINE AMP** or other Group 3 or 11 fungicides with different groups that control the same pathogens. Adhere to the label instructions regarding the consecutive use of **HEADLINE AMP** or other target site of action Group 3 or 11 fungicides that have a similar site of action on the same pathogens. To maintain the performance of **HEADLINE AMP** in the field, do not exceed the total number of applications of **HEADLINE AMP** per season as stated below in the Restrictions and Limitations table. Use tank mixtures with fungicides from a different group that is effective on the target pathogen when such use is permitted.
2. **IPM: HEADLINE AMP** should be integrated into an overall disease and pest management program. Use should be based on an IPM program that includes scouting, historical information related to pesticide use and crop rotation and considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
3. **Monitor:** Monitor efficacy of all fungicides used in the disease management program against the targeted pathogen and record other factors that may influence fungicide performance and/or disease development. If disease continues to progress after treatment with this product, do not increase the use rate. Discontinue use of this product, and switch to another fungicide with a different target site of action, if available.

4. **Reporting:** If a Group 3 or 11 target site fungicide, such as **HEADLINE AMP**, appears to be less or no longer effective against a pathogen that it previously controlled or suppressed, contact a BASF representative, local extension specialist, or certified crop advisor to assist in determining the cause of reduced performance.
5. Consult your local extension specialist, certified crop advisor and/or BASF for additional IPM strategies established for your area. **HEADLINE AMP** may be used in Agricultural Extension advisory (disease forecasting) programs, which recommend application timing based on environmental factors favorable for disease development. For further information and to report suspected resistance, contact a BASF representative at 1-877-371-2273 or at [www.agsolutions.ca](http://www.agsolutions.ca).

## MIXING

1. Clean spray tank following sprayer clean-up recommendations on the label of the product applied previously.
2. Fill the spray tank one-half full of water and start agitation.
3. Add the required amount of **HEADLINE AMP** to the tank.
4. Add the required amount of the tank mix partner, if required.
5. Continue agitation while filling the remainder of the spray tank.
6. After use, clean the spray tank.

## RESTRICTIONS AND LIMITATIONS

1. Do not enter or allow worker entry into treated areas during the restricted entry intervals specified in the following table:

Crop	Activity	Restricted Entry Interval (REI)
Corn	Hand harvesting, hand detasseling	13 days
	Hand set irrigation	1 day
	All other activities	12 hours
All other crops	All activities	12 hours

- Crop preharvest interval and application limitation:

Crop	Application to Harvest Interval (days)		Maximum Number of Applications per Year*
Wheat, barley, oats, rye and triticale	Apply no later than the end of flowering		2
Sweet corn	Mechanical harvesting	7	2
	Hand harvesting	13	
Field corn, popcorn and seed production corn	20		2

\*Do not make sequential applications of **HEADLINE AMP**.

- Rotational crops: A plant-back interval of 35 days is required for all crops not listed on the **HEADLINE AMP** label.

## PRECAUTIONS

- KEEP OUT OF REACH OF CHILDREN.**
- May be fatal or poisonous if swallowed. Do not take internally. Harmful if inhaled. Avoid inhaling/breathing dusts, sprays, etc.
- Causes moderate eye and skin irritation. DO NOT get in eyes or on skin.
- Wash thoroughly after handling and before eating, drinking or smoking.
- All mixer/loaders and applicators (except aerial applicators):** Wear coveralls over long sleeved shirt, long pants, chemical resistant gloves, socks and footwear during mixing/loading, application, clean-up and repair. In addition, wear goggles/face shield during mixing/loading, clean-up and repair. Gloves are not required during application.

**Aerial applicators:** Wear long sleeved shirt, long pants, socks and footwear during application.

Use closed cab tractor for ground boom application. A closed cab is a chemical resistant barrier that completely surrounds the occupant of the cab and prevents contact with the pesticide or treated surfaces outside the cab.

- If clothing becomes contaminated, remove and wash separately from household laundry before reuse.
- Clean spray equipment thoroughly after use.
- Apply only to agricultural crops when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

## FIRST AID

**If swallowed:** Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

## TOXICOLOGICAL INFORMATION

The patient should be treated symptomatically.

## ENVIRONMENTAL PRECAUTIONS

Toxic to aquatic organisms, non-target terrestrial plants and small wild mammals.

Observe buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g. sandy soil) and/or the depth to the water table is shallow.

## STORAGE

1. Store in original tightly closed container. Protect from freezing.
2. Do not ship or store near food, feed, seed and fertilizers.
3. Store in cool, dry, locked, well-ventilated area without floor drain.

## **DISPOSAL**

### **For Recyclable Containers**

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for any further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

### **For Non-Returnable Containers**

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

### **For Returnable-Refillable Containers**

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

## **NOTICE TO USER**

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.