

Veltyma DLX fungicide

technology sheet

Veltyma® DLX delivers protection against key leaf diseases and ear rots.

- Broad-spectrum management of both key leaf and ear diseases, including fusarium and gibberella ear rots
- Unique properties leading to better management of minor stress, better standability, increased growth efficiency and greener leaves¹
- Multiple modes of effective action for improved disease control and resistance management
- Reduced deoxynivalenol (DON) contamination to preserve grade quality

Active ingredients

Metconazole – Group 3
Mefentrifluconazole – Group 3
Pyraclostrobin– Group 11

Formulation

Caramba® – Liquid
Veltyma – Suspension concentrate

One case contains

1 x 8.1 L jug of Caramba
1 x 4.04 L jug of Veltyma

Storage

Requires heated storage.
Store away from food and feed.

Crops

Corn (field, pop, seed and sweet)

Timing

Full silking to silk browning

Diseases controlled

Common rust (*Puccinia sorghi*)
Eyespot (*Aureobasidium zeae*)
Fusarium ear rot (*Fusarium graminearum*)²
Gibberella ear rot (*Gibberella zeae*)²
Grey leaf spot (*Cercospora zeae-maydis*)
Northern corn leaf blight (*Setosphaeria turcica*)
Tar spot (*Phyllachora maydis*)

Broad-spectrum control of leaf and ear diseases



Untreated



Veltyma DLX

Source: BASF Small Plot Trials, Belmont, ON, 2021

¹ All comparisons are to untreated, unless otherwise stated.

² Suppression only.


 **BASF**

We create chemistry

Veltyma® DLX


Revysol® Fungicide

Improved reduction of gibberella




Source: BASF Small Plot Trials, inoculated, ON, 2021, n=3

Enhanced control of Northern corn leaf blight



Source: BASF Small Plot Trials, inoculated, ON, 2021, 9 weeks after application, n=20

Proven performance against tar spot



Source: BASF Small Plot Trials, ON, 2022, n=4

For more information:

Call **AgSolutions** Customer Care at 1-877-371-BASF (2273)

Visit agsolutions.ca

Application rates

One case of Veltyma DLX will treat 20 acres.

Caramba	405 ml/ac (1.0 L/ha)
Veltyma	202 ml/ac (500 ml/ha)

Water volume

Ground application	Minimum 80 L/ac (20 gal/ac)
Aerial application	20 L/ac (5 gal/ac)

Mixing order

1. Ensure the spray tank is clean before use. Follow the clean-out recommendations stated on the label of the product that was previously used.
2. Fill the spray tank 1/2 full of water and start agitation.
3. Add the required amount of Veltyma to the tank.
4. Add the required amount of Caramba to the spray tank.
5. Continue agitation while filling the remainder of the spray tank.
6. After use, clean the spray tank according to label precautions.

Application tips

Rainfastness – Do not apply when heavy rain is forecast.

Restricted entry interval – 12 hours.³

Resistance management – Fungicide use should be based on an integrated disease management program that includes scouting, historical information related to pesticide use and crop rotation and considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.

³ Except for hand harvesting corn (18 days) and hand set irrigation in corn (3 days).

Pre-harvest interval

21 days after application for corn.

Tank mixes

None on label.

Contact **AgSolutions**® Customer Care or your local BASF **AgSolutions** Retail Representative for additional information on supported tank mixes.

Always read and follow label directions.

AgSolutions, CARAMBA, REVYSOL and VELTYMA are registered trademarks of BASF; all used under license by BASF Canada Inc. CARAMBA, VELTYMA, and/or VELTYMA DLX fungicides should be used in a preventative disease control program.
© 2023 BASF Canada Inc.

Delaro and Proline are registered trademarks of Bayer Intellectual Property GmbH. Miravis is a registered trademark of Syngenta Participations AG.