

Veltyma fungicide

technology sheet

potatoes

The optimal fungicide for protection against key foliar diseases in potatoes, including early blight, black dot and brown spot.

- Multiple modes of effective action on early blight, including enhanced performance provided by the unique binding activity of Revysol®
- Proven **Plant Health Benefits**¹ for increased growth efficiency, better management of minor stress and greater yield potential²
- Delivers preventative and post-infection activity
- Liquid formulation for optimized usability

Active ingredients

Mefentrifluconazole – Group 3
Pyraclostrobin – Group 11

Formulation

Suspension concentrate


One case contains

2 x 8.1 L jugs

Storage

Store away from food and feed.

Potential yield increase with
Veltyma® fungicide



Source: Grower Applied Field Trial, NB, 2021

Crop

Potatoes

Timing

7 to 14 day application interval

Diseases controlled


In potatoes.

Black dot (*Colletotrichum coccodes*)

Brown spot (*Alternaria alternata*)³

Early blight (*Alternaria solani*)⁴

Potato yield compared to competitor



Source: Grower Applied Field Trials, PEI, NB, ON, AB, 2021-2022, n=9

¹ **Plant Health Benefits** refer to products that contain the active ingredient pyraclostrobin.

² All comparisons are to untreated unless otherwise stated.

³ Suppression.

⁴ Includes control of Group 11 resistant biotypes.

 **BASF**

We create chemistry

Application rate

One case of Veltyma will treat 32 hectares (80 acres).

Potatoes 500 ml/ha (202 ml/ac)

Mixing order

1. Ensure the spray tank is clean before use. Follow the clean-out recommendations stated on the label of the product that was previously used.
 2. Fill the spray tank 1/2 full of water and start agitation.
 3. Add the required amount of Veltyma to the tank.
 4. Add the required amount of the tank mix partner (if required).
 5. Continue agitation while filling the remainder of the spray tank.
 6. After use, clean the spray tank according to label precautions.
-

Application tips

Rainfastness – When product has dried on crop. Do not apply when heavy rain is forecasted.

Restricted entry interval – 12 hours.

Resistance management – Fungicide use should be based on an integrated disease management program that includes scouting, historical information related to pesticide use and crop rotation and considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.

Pre-harvest interval

7 days after application for potatoes.

Tank mixes

None on label.

Contact **AgSolutions®** Customer Care or your BASF **AgSolutions** Representative for information on supported tank mixes.

For more information

Call **AgSolutions** Customer Care at 1-877-371-BASF (2273) or visit agsolutions.ca.

Always read and follow label directions.

AgSolutions, REVYSOL and VELTYMA are registered trademarks of BASF; all used under license by BASF Canada Inc. VELTYMA fungicide should be used in a preventative disease control program. © 2024 BASF Canada Inc.