

Cantus fungicide

technology sheet

Proven protection against alternaria, botrytis and other key diseases in a wide range of crops.

- Rapidly rainfast to ensure long-lasting control under adverse conditions
- Protects against multiple fungal life stages, from initial infection to the spread of disease

Active ingredient

Boscalid – Group 7

Formulation


Wettable granules

One case contains


4 x 2.83 kg jugs

Storage

Store in cool, dry, ventilated area without floor drain.



4 applications made between 10% bloom and fruit set
Source: AAFC, Agassiz, BC, 2002



Source: BASF and university trials, 1997–2000

Crops

Berries

Bulb vegetables

Carrots

Cucurbit vegetables

Lettuce (field)

Fruiting vegetables

Grapes

Potatoes

Stone fruit

Strawberries

Diseases controlled

Botrytis gray mold (*Botrytis cinerea*)

Alternaria purple blotch (*Alternaria porri*)

Botrytis leaf blight (*Botrytis squamosa*)

Alternaria leaf blight (*Alternaria dauci*)

Alternaria blight (*Alternaria cucumerina*)

Gummy stem blight (*Didymella bryoniae*)

Botrytis rot (*Botrytis cinerea*)

Lettuce drop (*Sclerotinia minor*) – suppression

Botrytis gray mold (*Botrytis cinerea*)

Early blight (*Alternaria solani*)

Powdery mildew (*Uncinula necator*)

Early blight (*Alternaria solani*)

Brown rot (*Monilinia* spp.)

Monilinia blossom blight (*Monilinia* spp.)

Botrytis gray mold (*Botrytis cinerea*)

Timing

Refer to label for crop specific timing.

During periods of high disease pressure, use a shorter interval.

Application rates

One jug of Cantus treats 5 to 16.2 ha (12.3 to 40 acres).

Refer to label for crop specific rates and restrictions.

During periods of rapid growth or high disease pressure, use the higher rate.

Mixing order

1. Ensure the spray tank is clean before use.
 2. Fill the tank 1/2 full of water and start agitation.
 3. Add the required amount of Cantus fungicide to the tank.
 4. Continue agitation while filling the remainder of the spray tank.
 5. After use, clean the spray tank according to label precautions.
-

Application tips

Rainfastness – 1 hour.

Restricted entry interval – 12 hours.

Resistance management – Do not make more than two (2) sequential applications before alternating to a labeled fungicide with a different mode of action.

Pre-harvest interval

0 days for berries, carrots, cucurbits, fruiting vegetables, strawberries and stone fruit.

7 days for bulb vegetables.

14 days for field lettuce (head and leaf) and grapes.

30 days for potatoes.

Tank mixes

None on label.

Contact **AgSolutions**[®] Customer Care or your BASF Sales Representative for information on supported tank mixes.

For more information

Call **AgSolutions** Customer Care at 1-877-371-BASF (2273) or visit agsolutions.ca.



Always read and follow label directions.

AgSolutions is a registered trade-mark of BASF Corporation; CANTUS is a registered trade-mark of BASF SE; all used with permission by BASF Canada Inc. CANTUS fungicide should be used in a preventative disease control program. © 2015 BASF Canada Inc.