# Cantus fungicide technology sheet

# berries

## Proven, preventative protection against botrytis gray mold in berries.

- Rapidly rainfast to ensure long-lasting control under adverse conditions
- Locally systemic to protect treated areas, even as they expand
- Protects against multiple fungal life stages, from initial infection to the spread of disease

#### **Active ingredient**


Boscalid – Group 7

**Formulation** Wettable granules

**One case contains** 4 x 2.83 kg jugs

Storage

Store in cool, dry, ventilated area without floor drain.



4 applications made between 10% bloom and fruit set Source: AAFC, Agassiz, BC, 2002  $\,$ 

# Crop

Berries

## Timing

Early bloom, prior to onset of disease on a 7 to 14 day interval

During periods of high disease pressure, use a shorter interval.

## Disease controlled

Botrytis gray mold (Botrytis cinerea)





#### **Application rates**

#### One jug of Cantus fungicide treats 5.05 ha (12.5 acres).

Berries

560 g/ha (227 g/ac)

#### **Mixing order**

- 1. Ensure the spray tank is clean before use.
- 2. Fill the tank 1/2 full of water and start agitation.
- 3. Add the required amount of Cantus fungicide to the tank.
- 4. Continue agitation while filling the remainder of the spray tank.
- 5. After use, clean the spray tank according to label precautions.

#### Application tips

Rainfastness - 1 hour.

**Restricted entry interval** – 12 hours.

**Resistance management** – Do not make more than two (2) sequential applications before alternating to a labeled fungicide with a different mode of action.

#### **Pre-harvest interval**

0 days for berries.

#### Tank mixes

None on label.

Contact AgSolutions® Customer Care or your BASF Sales Representative for information on supported tank mixes.

#### For more information

Call AgSolutions Customer Care at 1-877-371-BASF (2273) or visit agsolutions.ca.

