

Cevya Revysol fungicide

technology sheet

Powered by Revysol®, an innovative new active ingredient, Cevya® fungicide is a systemic fungicide from BASF that provides fast and continuous pre- and post-infection control of key diseases.

- Fast and continuous control of key diseases in fruits and vegetables
- Preventative and post-infection control
- Unique, new binding activity to control biotypes that may have developed resistance to other Group 3, 7, 9 and 11 fungicides

Active ingredient

Mefentrifluconazole – Group 3

Formulation

Suspension concentrate


One case contains

2 x 4 L jugs

Storage


Store product away from food or feed.
Protect from freezing.

Grape powdery mildew control



Source: BASF, n=2

Apple scab incidence



Source: BASF

Crops

Grapes, sugar beets
Peanuts
Pome fruit
Potatoes, stone fruit, tree nuts

Timing

14 day application interval
10 to 14 day application interval
7 to 10 day application interval
7 to 14 day application interval

Cevya fungicide should be used preventatively.

Diseases controlled

In grapes.

Powdery mildew (*Erysiphe necator*)

In peanuts.

Early leaf spot (*Cercospora arachidicola*)¹

In pome fruit.

Apple scab (*Venturia inaequalis*)

Powdery mildew (*Podosphaera leucotricha*)¹

In potatoes.

Early blight (*Alternaria solani*)

Black dot (*Colletotrichum coccodes*)¹

Brown spot (*Alternaria alternata*)¹

In stone fruit.

Brown rot and blossom blight (*Monilinia fructicola*, *M. laxa*)¹

Powdery mildew (*Podosphaera clandestina*, *P. pannosa*)¹

In sugar beet.

Cercospora leaf spot (*Cercospora beticola*)

In tree nuts.

Brown rot and blossom blight (*Monilinia fructicola*, *M. laxa*)¹

Alternaria leaf spot (*Alternaria alternata*)¹

¹ Suppression.

 **BASF**

We create chemistry

Application rates

One jug of Cevya fungicide will treat 10.6 to 21 ha (26 to 52 acres).

Crops	Diseases	Application rates
Grapes	Powdery mildew	0.19 to 0.25 L/ha (0.075 to 0.1 L/ac)
Peanuts	Early leaf spot ¹	0.25 L/ha (0.1 L/ac)
Pome fruit	Apple scab, powdery mildew ¹	0.25 to 0.375 L/ha (0.1 to 0.15 L/ac)
Potatoes	Early blight Black dot ¹ Brown spot ¹	0.19 to 0.25 L/ha (0.075 to 0.1 L/ac)
Stone fruit	Brown rot and blossom blight ¹ , powdery mildew ¹	0.25 to 0.375 L/ha (0.1 to 0.15 L/ac)
Sugar beets	Cercospora leaf spot	0.19 to 0.375 L/ha (0.075 to 0.15 L/ac)
Tree nuts	Brown rot and blossom blight ¹ , alternaria leaf spot	0.25 to 0.375 L/ha (0.1 to 0.15 L/ac)

Under high disease pressure and during rapid growth, use the higher rate and shorter spray interval.

¹ Suppression.

Mixing order

1. Ensure the spray tank is clean before use.
2. Fill the spray tank 1/2 full of water and start agitation.
3. Add the required amount of Cevya to the tank.
4. Add the required amount of tank-mix partner, if applicable.
5. Add the recommended amount of adjuvant, if applicable.
6. Continue agitation while filling the remainder of the spray tank.
7. After use, clean the spray tank according to label precautions.

Application tips

Rainfastness

1 hour.

Restricted entry interval

12 hours for all crops and activities except girdling and cane turning in grapes (35 days).

Resistance management

Cevya is an excellent resistance management tool to include in an IPM program. It can be used in combination or rotation with other chemistries to prevent the development of resistant strains. To limit the potential for development of resistance, rotate the use of Cevya or other Group 3 fungicides with different groups that control the same pathogens.

Pre-harvest interval

Refer to label for crop-specific pre-harvest intervals.

Tank mixes

None on label.

Contact **AgSolutions**[®] Customer Care or your BASF Sales Representative for information on supported tank mixes.

For more information

Call **AgSolutions** Customer Care at 1-877-371-BASF (2273) or visit agsolutions.ca/horticulture.

Always read and follow label directions.

AgSolutions, CEVYA, and REVYSOL are registered trade-marks of BASF, used under license by BASF Canada Inc. CEVYA fungicide should be used in a preventative disease control program. © 2021 BASF Canada Inc.

INSPIRE SUPER[®] is a registered trademark of a Syngenta Group Company. Luna Tranquility[®] is a registered trademark of the Bayer Group.