

# Cevya®

Revysol® Fungicide

Cevya® fungicide is powered by Revysol® to provide fast, systemic, continuous pre- and post-infection control of key diseases.

- Fast and continuous control of key diseases in potatoes
- Preventative and post-infection control
- Unique, new binding activity to control biotypes that may have developed resistance to other Group 3, 7, 9 and 11 fungicides
- Timing and tank-mix flexibility to adapt to the season's needs

## Active ingredient

Mefentrifluconazole – Group 3

## Formulation

Suspension concentrate

## One case contains

2 x 4 L jugs

## Storage

Requires heated storage.

## Crops

Potatoes

## Timing

7 to 14 day application interval

Cevya fungicide should be used preventatively.

## Diseases controlled


### In potatoes.

Early blight (*Alternaria solani*)

Black dot (*Colletotrichum coccodes*)<sup>1</sup>

Brown spot (*Alternaria alternata*)<sup>1</sup>

### Early blight control



Source: BASF Small Plot Trials, 2016-2018, n=3



<sup>1</sup> Suppression.

 **BASF**

We create chemistry

## Application rates

**One case of Cevya fungicide will treat 32 to 42 ha (80 to 104 acres).**

Potatoes 0.19 to 0.25 L/ha (0.075 to 0.1 L/ac)

Under high disease pressure and during rapid growth, use the higher rate and shorter spray interval.

---

## Mixing order

1. Ensure the spray tank is clean before use.
  2. Fill the spray tank 1/2 full of water and start agitation.
  3. Add the required amount of Cevya to the tank.
  4. Add the required amount of tank-mix partner, if applicable.
  5. Continue agitation while filling the remainder of the spray tank.
  6. After use, clean the spray tank according to label precautions.
- 

## Application tips

**Rainfastness** - 1 hour.

**Restricted entry interval** - 12 hours.

**Resistance management** - Cevya is an excellent resistance management tool to include in an IPM program. It can be used in combination or rotation with other chemistries to prevent the development of resistant strains. To limit the potential for development of resistance, rotate the use of Cevya or other Group 3 fungicides with different groups that control the same pathogens.

---

## Pre-harvest interval

7 days.

---

## Tank mixes

None on label.

Contact your local BASF **AgSolutions**<sup>®</sup> Grower or Retail Representative or call **AgSolutions** Customer Care at 1-877-371-BASF (2273) for information on supported tank mixes.

---

**For more information:** Call **AgSolutions** Customer Care at 1-877-371-BASF (2273) or visit [agsolutions.ca](https://agsolutions.ca).

**Always read and follow label directions.**