Forum fungicide technology sheet

Highly systemic tank-mix partner for downy mildew control in grapes.

- Antisporulant activity controls spores and stops the spread of disease
- Easy-to-use liquid formulation

Active ingredient Dimethomorph – Group 40

Formulation Flowable concentrate

One case contains 2 x 4.5 L jugs

Storage Store in cool, dry, ventilated area away from food and animal feed.



Crop

Grapes

Timing

Begin prior to onset of disease on a 7 to 10-day interval

During periods of rapid growth or high disease pressure, use a shorter interval.

Disease controlled

Downy mildew (Plasmopara viticola)



Downy mildew control in grapes

Source: Summary of 6 BASF & 3rd party trials





Application rates

One jug of Forum treats 10 ha (25 acres).

Grapes

450 mL/ha (182 mL/ac)

Mixing order

1. Ensure the spray tank is clean before use.

- 2. Fill the tank 1/2 full of water and start agitation.
- 3. Add the required amount of Forum fungicide to the tank while agitating.
- 4. Add the required amount of tank-mix partner.
- 5. Continue agitation while filling the remainder of the spray tank.
- 6. After use, clean the spray tank according to label precautions.

Application tips

Rainfastness – 2 hours.

Restricted entry interval – 12 hours.

Do not enter treated areas for cane turning or girdling within 19 days after application, or for training, thinning or hand pruning within 12 days after application.

Resistance management – In order to reduce the risk of developing fungicide resistance, Forum fungicide must be used in a tank mix with other fungicides effective against downy mildew.

Pre-harvest interval

14 days for grapes.

Tank mixes

None on label.

Contact **AgSolutions**[®] Customer Care or your BASF Sales Representative for additional information on supported tank mixes.

For more information

Call AgSolutions Customer Care at 1-877-371-BASF (2273) or visit agsolutions.ca/horticulture.

Always read and follow label directions.