

# Forum®

## Fungicide

### Excellent activity on late blight in potatoes, both in the field and into storage.

- Highly systemic tank-mix partner for control of late blight in potatoes
- Antisporulant activity controls spores and stops the spread of disease
- Easy-to-use liquid formulation

#### Active ingredient

Dimethomorph – Group 40

#### Formulation

Suspension concentrate

#### One case contains

2 x 4.5 L jugs

#### Storage

Requires heated storage.

#### Crop<sup>1</sup>

Potatoes

#### Timing

5 to 10 day interval


During periods of rapid growth or high disease pressure, use a shorter interval. See label for details.

#### Diseases controlled

Late blight (*Phytophthora infestans*)


Suppression of tuber blight in storage (*Phytophthora infestans*)

Curative late blight control



Source: BASF Small Plot Trials, 2001-2006, n=17

Storage rot control with Forum fungicide and protectant applied pre-harvest



Source: BASF Small Plot Trials, 2001-2006, n=7



<sup>1</sup> Refer to label for additional crops.

 **BASF**

We create chemistry

## Application rates

**One case of Forum fungicide will treat 50 acres (20.2 ha).**

Potatoes 182 ml/ac (450 ml/ha)

Always apply Forum as a tank mix with a protectant/contact fungicide to protect against late blight and for resistance management.

---

## Mixing order

1. Ensure the spray tank is clean before use.
  2. Fill the tank 1/2 full of water and start agitation.
  3. Add the required amount of Forum fungicide to the tank while agitating.
  4. Add the required amount of fungicide tank mix partner to the tank.
  5. Continue agitation while filling the remainder of the spray tank.
  6. After use, clean the spray tank according to label precautions.
- 

## Application tips

Apply a maximum of three applications per season.

**Rainfastness** – 2 hours.

**Restricted entry interval** – 12 hours.

**Resistance management** – In order to reduce the risk of developing fungicide resistance, Forum fungicide must be used in a tank mix with other fungicides effective against late blight.

---

## Pre-harvest interval

4 days after application for potatoes.

---

## Tank mixes

**Fungicides:** See label for details.

Contact your local BASF **AgSolutions**<sup>®</sup> Grower or Retail Representative or call **AgSolutions** Customer Care at 1-877-371-BASF (2273) for information on supported tank mixes.

---

**For more information:** Call **AgSolutions** Customer Care at 1-877-371-BASF (2273) or visit [agsolutions.ca](https://agsolutions.ca).

**Always read and follow label directions.**

AgSolutions and FORUM are registered trademarks of BASF; all used under license by BASF Canada Inc. © 2024 BASF Canada Inc.