

# Headline fungicide

## technology sheet

Headline® fungicide provides systemic control of both early blight and late blight, the two most devastating diseases of potatoes.

- Proven, effective control of early and late blight, including sensitive strains
- Will not wash off; it's systemic and rainfast in under 1 hour to ensure long-lasting control, even under adverse conditions
- Excellent tank-mix or rotation partner with fungicides of a different mode of action

### Active ingredient

Pyraclostrobin – Group 11

### Formulation

Emulsifiable concentrate

### One case contains

2 x 6.5 L jugs

### Storage

Store product away from food, feed, seed and fertilizers.

Protect from freezing.

### Crops

Potatoes

### Timing

Prior to row closure or when conditions become favourable for the development of disease (whichever comes first).

Headline fungicide should be used preventatively.


### Diseases controlled

#### In potatoes.

Early blight (*Alternaria solani*)

Late blight (*Phytophthora infestans*)

Headline at Tuber Initiation Yield Trial



Source: Felix Farms

**BASF**  
We create chemistry

**AgCelence®**  
Expect more

## Application rates

One jug will treat 9.7 to 14.4 ha (24 to 35.7 acres).

Crops	Diseases	Application Rates
Potatoes	Early blight	0.45 to 0.67 L/ha (0.18 to 0.27 L/ac)
	Late blight	0.45 to 0.67 L/ha (0.18 to 0.27 L/ac)

Under high disease pressure and during rapid growth, use the higher rate and tank mix it with a multisite fungicide such as Bravo® 500 Agricultural Fungicide. Refer to the respective tank-mix partner label for rates, additional recommendations, restrictions and precautions.

## Mixing order

1. Ensure the spray tank is clean before use.
2. Fill the spray tank one-half full of water and start agitation.
3. Add the required amount of Headline to the tank.
4. Add the required amount of tank-mix partner, if applicable.
5. Add the recommended amount of adjuvant, if applicable.
6. Continue agitation while filling the remainder of the spray tank.
7. After use, clean the spray tank according to label precautions.

## Application tips

**Rainfastness** – 1 hour

**Restricted entry interval** – 12 hours

**Resistance management** – To limit the potential for development of resistance DO NOT apply more than one (1) application before rotating to another mode of action for at least one application. No more than three (3) applications should be made per season.

## Pre-harvest interval

6 days

## Tank mixes

Contact **AgSolutions®** Customer Care or your BASF Sales Representative for information on supported tank mixes.

## For more information:

Call **AgSolutions** Customer Care at 1-877-371-BASF (2273) or visit [agsolutions.ca](http://agsolutions.ca).

## Always read and follow label directions.

**AgSolutions**, and **HEADLINE** are registered trade-marks of BASF. **HEADLINE** fungicide should be used in a preventative disease control program. © 2020 BASF Canada Inc. **BRAVO®** is a registered trademark of a Syngenta Group Company.