

Pristine fungicide technology sheet

berries

Preventative, broad-spectrum protection with the benefits of AgCelence[®].

- Multiple modes of action for broad-spectrum protection against key diseases, including anthracnose and botrytis
- Unique **AgCelence** benefits¹ for improved stress management and increased yield potential
- Locally systemic and rapidly rainfast

Active ingredients

Boscalid – Group 7
Pyraclostrobin – Group 11

Formulation

Wettable granules

One case contains

4 x 2.83 kg jugs

Storage

Store in cool, dry, well-ventilated area without floor drain.

Crops

Blackberries, blueberries, raspberries and strawberries²

Timing

5 to 10% bloom or prior to onset of disease on a 7 to 14 day interval

Additional crops on label: Currants, elderberries, gooseberries, huckleberries, loganberries and Saskatoon berries.

During periods of rapid growth or high disease pressure, use a shorter interval.


Diseases controlled

In blackberries, blueberries and raspberries.

Anthrachnose (*Colletotrichum* spp.)
Botrytis gray mold (*Botrytis cinerea*)
Phomopsis
Septoria leaf spot (*Septoria albopunctata*)³
Valdensinia leaf spot (*Valdensinia heterodoxa*)³

In strawberries.

Anthrachnose (*Colletotrichum* spp.)
Botrytis gray mold (*Botrytis cinerea*)
Leaf spot (*Mycosphaerella* spp.)
Powdery mildew (*Sphaerotheca* spp.)



Source: BASF & Independent 3rd party research

¹ AgCelence benefits refer to products that contain the active ingredient pyraclostrobin. All comparisons are against grower standard unless otherwise stated.

² Not for use in strawberry nurseries.

³ Suppression in lowbush blueberries grown in Eastern Canada only. Not controlled in other crops.

Application rates

One jug of Pristine® will treat 1.8 to 2.2 ha (4.4 to 5.4 acres).

Crops	Diseases	Application rates
Blackberries, raspberries	Anthracnose, botrytis gray mold	1.3 to 1.6 kg/ha (526 to 647 g/ac)
	Phomopsis	1.6 kg/ha (647 g/ac)
Blueberries	Anthracnose, botrytis gray mold, valdensinia leaf spot	1.3 to 1.6 kg/ha (526 to 647 g/ac)
	Phomopsis, septoria leaf spot	1.6 kg/ha (647 g/ac)
Strawberries	Anthracnose, botrytis gray mold, leaf spot	1.3 to 1.6 kg/ha (526 to 647 g/ac)
	Powdery mildew	1.6 kg/ha (647 g/ac)

During periods of rapid growth or high disease pressure, use the higher rate.

Mixing order

1. Ensure the spray tank is clean before use.
2. Fill the tank 1/2 full of water and start agitation.
3. Add the required amount of Pristine fungicide to the tank.
4. Continue agitation while filling the remainder of the spray tank.
5. After use, clean the spray tank according to label precautions.

Application tips

Rainfastness – 1 hour.

Restricted entry interval – Once dry to 24 hours. Refer to label for details.

Resistance management – Do not apply more than two (2) sequential applications of Pristine fungicide before alternating to a fungicide with a different mode of action other than Group 7 and Group 11. Group 11-containing fungicide applications should be no more than one-half of the total number of fungicide applications per season.

Pre-harvest interval

1 day for strawberries.

0 days for all other registered berry crops.

Tank mixes

None on label.

For more information

Call **AgSolutions®** Customer Care at 1-877-371-BASF (2273) or visit agsolutions.ca.

Always read and follow label directions.

AgSolutions is a registered trade-mark of BASF Corporation; **AgCelence**, and **PRISTINE** are registered trade-marks of BASF SE; all used with permission by BASF Canada Inc. PRISTINE fungicide should be used in a preventative disease control program. © 2015 BASF Canada Inc.