

Pristine fungicide

technology sheet

hops

Preventative, broad-spectrum protection with the benefits of AgCelence[®].

- Multiple modes of action for control of powdery mildew and suppression of downy mildew
- Targets multiple stages of the disease cycle to prevent initial infection and disease spread
- Locally systemic and rapidly rainfast
- Unique **AgCelence** benefits¹ for improved stress management and increased yield potential

Active ingredients

Boscalid – Group 7
Pyraclostrobin – Group 11

Formulation

Wettable granules

One case contains

4 x 2.83 kg jugs

Storage

Store in cool, dry, well-ventilated area without floor drain.

Crops

Hops

Timing

10 to 14 day interval,
max. 3 applications per season

Apply preventatively prior to disease development.
During periods of rapid growth or high disease pressure, use a shorter interval.

Diseases controlled

In hops.

Downy mildew (*Pseudoperonospora humuli*)²
Powdery mildew (*Podosphaera macularis*)

¹ **AgCelence** benefits refer to products that contain the active ingredient pyraclostrobin. All comparisons are against grower standard unless otherwise stated.

² Suppression only.

Application rates

One jug of Pristine® will treat 1.8 to 11.3 ha (4.4 to 27 acres).

Hops 105 g per 100 L dilute spray, max. 1524 L/ha

Use 234 to 468 L of dilute spray per hectare (0.25 to 0.49 kg/ha) prior to trellising and 935 to 1524 L of dilute spray per hectare (0.98 to 1.6 kg/ha) thereafter.

Mixing order

1. Ensure the spray tank is clean before use.
 2. Fill the tank 1/2 full of water and start agitation.
 3. Add the required amount of Pristine fungicide to the tank.
 4. Continue agitation while filling the remainder of the spray tank.
 5. After use, clean the spray tank according to label precautions.
-

Application tips

Rainfastness – 1 hour.

Restricted entry interval – 12 hours to 46 days.

Refer to label for details.

Resistance management – Do not apply more than two (2) sequential applications of Pristine fungicide before alternating to a fungicide with a mode of action other than Group 7 and Group 11. Maximum three (3) applications of Pristine fungicide per season.

Pre-harvest interval

46 days for hops.

Tank mixes

None on label.

For more information

Call **AgSolutions®** Customer Care at 1-877-371-BASF (2273) or visit **agsolutions.ca**.

Always read and follow label directions.

AgSolutions is a registered trade-mark of BASF Corporation; AgCelence, and PRISTINE are registered trade-marks of BASF SE; all used with permission by BASF Canada Inc. PRISTINE fungicide should be used in a preventative disease control program. © 2015 BASF Canada Inc.