# Sercadis fungicide

# technology sheet

Innovative, new chemistry for consistent, continuous control of key diseases.

- Control of scab and powdery mildew in pome fruit
- Tank-mix and rotational flexibility to adapt to the season's needs
- Highly systemic activity helps protect new growth

# **Active ingredient**

Fluxapyroxad – Group 7

### **Formulation**

Suspension

#### One case contains

2 x 1.35 L jugs

#### **Storage**

Store away from food or feed.



# **Crops**

Pome fruits

#### **Timing**

Begin prior to onset of disease on a 7 to 14 day interval

#### **Diseases controlled**

### In pome fruits.

Apple scab (Venturia inaequalis)

Pear scab (Venturia pirina)

Powdery mildew (Podosphaera leucotricha)





### **Application rates**

#### One jug of Sercadis™ fungicide treats 4 to 8 ha (10 to 20 acres).

Pome fruits

Apple scab 333 mL/ha (135 mL/ac)

Powdery mildew 167 to 333 mL/ha (67 to 135 mL/ac)

#### Mixing order

- 1. Ensure the spray tank is clean before use.
- 2. Fill the tank 1/2 full of water and start agitation.
- 3. Add the required amount of Sercadis fungicide to the tank.
- 4. Add the required amount of tank mix partner, if applicable.
- 5. Add the recommended amount of adjuvant, if applicable.
- 6. Continue agitation while filling the remainder of the spray tank.
- 7. After use, clean the spray tank according to label precautions.

## **Application tips**

Rainfastness - 1 hour.

Restricted entry interval – 12 hours.

**Resistance management** – Tank-mix with a non-Group 7 fungicide when such use is permitted. Avoid consecutive applications of Sercadis or other Group 7 fungicides in a season.

Use of a non-ionic surfactant at 0.125% v/v is recommended.

#### Pre-harvest interval

0 days for pome fruits.

### Tank mixes

None on label.

Contact AgSolutions® Customer Care or your BASF Sales Representative for information on supported tank mixes.

#### For more information

Call AgSolutions Customer Care at 1-877-371-BASF (2273) or visit agsolutions.ca.