



We create chemistry

**PRODUCT LABELS FOR
ALTITUDE FX3:**

**AC 299,263 120 AS
HERBICIDE SOLUTION**

+

STARANE II HERBICIDE

| | | |
|-------|---|-----------|
| GROUP | 2 | HERBICIDE |
|-------|---|-----------|

AC 299,263 120 AS

Herbicide Solution

FOR SALE FOR USE IN THE PRAIRIE PROVINCES AND INTERIOR OF BRITISH COLUMBIA
(INCLUDING THE PEACE RIVER REGION) ONLY ON **Clearfield®** AND **Clearfield® Plus**
WHEAT¹ (WHEAT VARIETIES WITH THE **Clearfield** TRAIT) AND FIELD PEAS

COMMERCIAL (AGRICULTURAL)

ACTIVE INGREDIENT: Imazamox120 g/L (present as ammonium salt)

Contains 1,2 benzisothiazolin-3-one at 0.019% as a preservative

REGISTRATION NO. 26705 PEST CONTROL PRODUCTS ACT

NET CONTENTS: 1.34 L – 200 L

IN CASE OF EMERGENCY ENDANGERING LIFE OR PROPERTY
INVOLVING THIS PRODUCT, CALL DAY OR NIGHT
1-800-454-2673

READ THE LABEL AND THE BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

BASF Canada Inc.
5025 Creekbank Road
Building A, 2nd Floor
Mississauga, Ontario L4W 0B6
1-877-371-2273

Clearfield is a registered trade-mark of BASF Agrochemical Products B.V., used under license permission by BASF Canada Inc.

(Back Panel of the Jug)

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

DO NOT APPLY BY AIR.

Harmful if absorbed through skin or inhaled. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing.

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. In addition, wear goggles or face shield for mixing/loading, clean-up and repair activities. Gloves are not required during application within a closed cab.

Do not enter or allow workers entry into treated areas during the restricted entry interval (REI) of 12 hours.

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

ENVIRONMENTAL PRECAUTIONS

TOXIC to non-target terrestrial plants. Observe spray buffer zones specified under Directions for Use.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g. sandy soil) and/or the depth to the water table is shallow.

FIRST AID

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

STORAGE

Store above 5°C.

Store in a cool, dry place away from children, animals, food, feed or fertilizers.

Keep from freezing.

DISPOSAL

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

¹ *Methods of growing plants which are resistant to certain acetohydroxyacid synthase (AHAS) herbicides are protected by Canadian Patent No. 1341465. Other applicable patents include Canadian Patent Nos. 2456314, 2455512, 2456311, 2492167 and 2790951. The use of this product in the practice of patented methods could constitute patent infringement. Further information can be obtained by contacting BASF Canada at 1-877-371-2273.*

| | | |
|-------|---|-----------|
| GROUP | 2 | HERBICIDE |
|-------|---|-----------|

AC 299,263 120 AS

Herbicide Solution

FOR SALE FOR USE IN THE PRAIRIE PROVINCES AND INTERIOR OF BRITISH COLUMBIA
(INCLUDING THE PEACE RIVER REGION) ONLY ON **Clearfield®** AND **Clearfield® Plus**
WHEAT¹ (WHEAT VARIETIES WITH THE **Clearfield** TRAIT) AND FIELD PEAS

COMMERCIAL (AGRICULTURAL)

ACTIVE INGREDIENT: Imazamox120 g/L (present as ammonium salt)

Contains 1,2 benzisothiazolin-3-one at 0.019% as a preservative

REGISTRATION NO. 26705

PEST CONTROL PRODUCTS ACT

NET CONTENTS: 1.34 L – 200 L

IN CASE OF EMERGENCY ENDANGERING LIFE OR PROPERTY
INVOLVING THIS PRODUCT, CALL DAY OR NIGHT
1-800-454-2673

READ THE LABEL AND THE BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

BASF Canada Inc.
5025 Creekbank Road
Building A, 2nd Floor
Mississauga, Ontario L4W 0B6
1-877-371-2273

Clearfield is a registered trade-mark of BASF Agrochemical Products B.V., used under license
by BASF Canada Inc.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

DO NOT APPLY BY AIR.

Harmful if absorbed through skin or inhaled. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing.

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. In addition, wear goggles or face shield for mixing/loading, clean-up and repair activities. Gloves are not required during application within a closed cab.

Do not enter or allow workers entry into treated areas during the restricted entry interval (REI) of 12 hours.

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

ENVIRONMENTAL PRECAUTIONS

TOXIC to non-target terrestrial plants. Observe spray buffer zones specified under Directions for Use.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g. sandy soil) and/or the depth to the water table is shallow.

FIRST AID

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

STORAGE

Store above 5°C.

Store in a cool, dry place away from children, animals, food, feed or fertilizers.

Keep from freezing.

DISPOSAL

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

DIRECTIONS FOR USE (Prairie Provinces and Interior of British Columbia)

AC 299,263 120 AS Herbicide Solution is a selective herbicide that can be applied as an early post-emergent treatment in **Clearfield®** or **Clearfield® Plus** Wheat¹ varieties (wheat varieties with the **Clearfield** trait). **Clearfield** and **Clearfield Plus** Wheat¹ varieties are tolerant of imazamox, the active ingredient in AC 299,263 120 AS Herbicide Solution. Wheat varieties that are not designated as having the **Clearfield** trait may be damaged or destroyed by this treatment. When AC 299,263 120 AS Herbicide Solution is applied early post-emergence, absorption may occur through both the roots and foliage. Susceptible weeds stop growing and eventually die.

AC 299,263 120 AS Herbicide Solution may also be applied as an early post-emergence treatment in field peas when applied in tank mix with Basagran Forte Herbicide. **DO NOT** apply to field peas that have been subjected to stress from conditions such as hail damage, flooding, drought, hot, humid weather, widely fluctuating temperature conditions, prolonged cold weather or injury from prior herbicide applications, as crop injury may result.

Apply using ground equipment only. **DO NOT** apply by air.

Do not apply more than once per year.

DO NOT apply directly to aquatic habitats (such as lakes, rivers, sloughs, ponds, coulees, prairie potholes, creeks, marshes, streams, reservoirs, ditches and wetlands) and estuarine/marine habitats.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

As this product is not registered for the control of pests in aquatic systems, **DO NOT** use to control aquatic pests.

DO NOT apply during periods of dead calm or when winds are gusty.

DO NOT graze treated wheat within 14 days of application or cut for hay within 42 days of application.

DO NOT graze treated field peas or cut for hay within 20 days of application.

Pre-harvest Interval: Wheat grain and straw can be harvested 79 days after treatment. After 60 days, field peas can be harvested.

Field Sprayer Application

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Boom height must be 60 cm or less above the crop or ground.

SPRAY BUFFER ZONES

A spray buffer zone is NOT required for:

- Uses with hand-held application equipment permitted on this label.
- Low-clearance hooded or shielded sprayers that prevent spray contact with crop, fruit or foliage.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands).

| Method of Application | Crop | Spray Buffer zones (metres) Required for the Protection of Terrestrial Habitat |
|-----------------------|------------------------------|--|
| Field sprayer | Wheat and peas (dried field) | 1 |

For tank mixes, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pest Management Regulatory Agency web site.

CROP: Clearfield® and Clearfield® Plus Wheat¹ (Wheat varieties with the Clearfield trait)

BROADLEAF WEED AND GRASS CONTROL

Application should be made from the 2 to the 6 leaf stage of wheat crop and after the weeds have emerged. To control broadleaf weeds apply AC 299,263 120 AS Herbicide Solution from the cotyledon up to the 4 leaf stage. To control grasses, apply AC 299,263 120 AS Herbicide Solution from the 1-4 true leaf stage up until early tillering. Initial crop injury may be observed after application but this is outgrown and should not affect yield. Severe crop injury will occur as a result of spray overlap. AVOID SPRAYER OVERLAP.

| | |
|---|---|
| AC 299,263 120 AS Herbicide Solution at 125 mL product/ha (15 g a.e./ha) will control: | |
| Lamb's-quarters* Stinkweed Wild mustard | Volunteer spring wheat (non- Clearfield wheat) |
| AC 299,263 120 AS Herbicide Solution at 167 mL product/ha (20 g a.e./ha) will control: | |
| Barnyard grass Cleavers* Cow cockle Green foxtail Green smartweed Japanese Brome Grass* Lamb's-quarters* Persian damel Redroot pigweed Round-leaved mallow* Russian thistle* Shepherd's purse Stinkweed | Volunteer barley Volunteer canary seed Volunteer canola (non- Clearfield canola only) Volunteer durum wheat Volunteer spring wheat (non- Clearfield wheat) Volunteer tame oats Wild buckwheat* Wild mustard Wild oats Yellow foxtail |

* = suppression only

APPLICATION INSTRUCTIONS

Apply in 50-100 L/ha of water. Applications of AC 299,263 120 AS Herbicide Solution require the addition of an adjuvant as indicated in the table below. For **Clearfield** wheat varieties, apply AC 299,263 120 AS Herbicide Solution with a non-ionic surfactant (NIS) only. For **Clearfield Plus** wheat varieties, AC 299,263 120 AS Herbicide Solution may be applied with MSO Concentrate with Leci-Tech, Merge Adjuvant or NIS. Refer to the **Adjuvant Options** table below for rates.

Adjuvant Options

| Wheat Variety | Non-Ionic Surfactant ¹ | MSO Concentrate with Leci-Tech | Merge Adjuvant |
|------------------------|-----------------------------------|--------------------------------|----------------|
| Clearfield | 0.25% v/v | — | — |
| Clearfield Plus | 0.25% v/v | 1% v/v | 0.5% v/v |

¹ Examples of non-ionic surfactants (NIS) include AGRAL® 90, AG-SURF® or SURF® 92.

DO NOT use MSO or Merge Adjuvant with AC 299,263 120 AS Herbicide Solution on **Clearfield** wheat varieties. MSO or Merge Adjuvant may be used with AC 299,263 120 AS Herbicide Solution on **Clearfield Plus** only wheat varieties. When AC 299,263 120 AS Herbicide Solution is tank mixed with another herbicide, use of MSO or Merge Adjuvant on **Clearfield Plus** wheat varieties is only recommended when a tank mix partner allows use of MSO or Merge Adjuvant.

TANK-MIXING

To provide broad-spectrum weed control in **Clearfield** and **Clearfield Plus** wheat varieties, AC 299,263 120 AS Herbicide Solution may be tank-mixed with the following tank-mix partners. When applied as a tank-mix combination, read and observe all label directions, including rates, personal protective equipment, restrictions and precautions for each product used in the tank-mix. Always use in accordance with the most restrictive label restrictions and precautions.

| AC 299,263 120 AS Herbicide Solution + 2,4-D Ethylhexyl Ester | | |
|--|--|----------------------------------|
| Crops | Clearfield and Clearfield Plus Wheat ¹ | |
| Timing | Post-emergent application from the 2 to 6-leaf stage of crop | |
| Tank Mix Components | AC 299,263 120 AS Herbicide Solution | 125 mL/ha OR 167 mL/ha |
| | 2,4-D Ester | 560 g ai/ha |
| Adjuvant | Refer to table of Adjuvant Options above. Adjuvant option and rate will depend on wheat variety. | |
| Spray Volume | 50-100 L/ha water | |
| Weeds Controlled | This tank-mix combination controls all weeds as listed on the AC 299,263 120 AS Herbicide Solution and 2,4-D Ester herbicide labels. | |

| AC 299,263 120 AS Herbicide Solution + CURTAIL® M Herbicide | | |
|--|--|---|
| Crops | Clearfield and Clearfield Plus Wheat ¹ | |
| Timing | Post-emergent application from the 3 to 6-leaf stage of crop | |
| Tank Mix Components | AC 299,263 120 AS Herbicide Solution | 125 mL/ha OR 167 mL/ha |
| | CURTAIL M | A maximum rate of 660 g ai/ha (2.0 L pdt/ha) |
| Adjuvant | Refer to table of Adjuvant Options above. Adjuvant option and rate will depend on wheat variety. | |
| Spray Volume | 50-100 L/ha water | |
| Weeds Controlled | This tank-mix combination controls all weeds as listed on the AC 299,263 120 AS Herbicide Solution and Curtail M Herbicide labels. | |

| AC 299,263 120 AS Herbicide Solution + FRONTLINE® Herbicide | | |
|--|--|----------------------------------|
| Crops | Clearfield and Clearfield Plus Wheat¹ | |
| Timing | Post-emergent application from the extended 3 to 6 leaf stage of crop. | |
| Tank Mix Components | AC 299,263 120 AS Herbicide Solution | 125 mL/ha OR 167 mL/ha |
| | FRONTLINE A | 0.10 L/ha |
| | FRONTLINE B | 0.840 L/ha |
| Adjuvant | Refer to table of Adjuvant Options above. Adjuvant option and rate will depend on wheat variety. | |
| Spray Volume | 50-100 L/ha water | |
| Weeds Controlled | This tank-mix combination controls all weeds as listed on the AC 299,263 120 AS Herbicide Solution and Frontline Herbicide labels. | |

| AC 299,263 120 AS Herbicide Solution + Starane™ or Starane II Herbicide | | |
|--|---|--|
| Crops | Clearfield and Clearfield Plus Wheat¹ | |
| Timing | Post-emergent application from the 2 to 6 true leaf stage of crop | |
| Tank Mix Components | AC 299,263 120 AS Herbicide Solution | 167 mL/ha |
| | Starane or Starane II Herbicide | 0.6 L/ha Starane OR 0.31 L/ha Starane II |
| Adjuvant | Refer to table of Adjuvant Options above. Adjuvant option and rate will depend on wheat variety. | |
| Spray Volume | 50-100 L/ha water | |
| Weeds Controlled | In addition to the control of weeds as listed on the AC 299,263 120 AS Herbicide Solution label, this tank-mix combination controls kochia (including Group 2 resistant), volunteer flax and cleavers (including Group 2 resistant), and provides suppression of stork's bill. Initial crop injury may be observed after application but this is outgrown and should not affect yield. Severe crop injury and a delay in maturity may occur as a result of over spray. AVOID SPRAYER OVERLAP. | |

| AC 299,263 120 AS Herbicide Solution + Starane™ or Starane II Herbicide + MCPA Ester | | |
|---|---|---|
| Crops | Clearfield and Clearfield Plus Wheat¹ | |
| Timing | Post-emergent application from the 3 to 6 true leaf stage of crop | |
| Tank Mix Components | AC 299,263 120 AS Herbicide Solution | 167 mL/ha |
| | Starane or Starane II Herbicide | 0.6 L/ha Starane OR 0.31 L/ha Starane II |
| | MCPA Ester | 560 g ai/ha (500 g/L formulation: 1.12 L/ha) (600 g/L formulation: 0.93 L/ha) |
| Adjuvant | Refer to table of Adjuvant Options above. Adjuvant option and rate will depend on wheat variety. | |
| Spray Volume | 50-100 L/ha water | |
| Weeds Controlled | <p>This tank-mix combination controls cow cockle, wild buckwheat and common chickweed (non-Group 2 resistant biotypes) in addition to all weeds as listed on the AC 299,263 120 AS Herbicide Solution label plus weeds as listed on the Starane Herbicide label for the tank-mix of Starane Herbicide plus MCPA Ester.</p> <p>Initial crop injury may be observed after application but this is outgrown and should not affect yield. Severe crop injury and a delay in maturity may occur as a result of over spray. AVOID SPRAYER OVERLAP.</p> | |

| AC 299,263 120 AS Herbicide Solution + Starane™ or Starane II Herbicide + 2,4-D Ester Herbicide | | |
|--|--|--|
| Crops | Clearfield and Clearfield Plus Wheat¹ | |
| Timing | Post-emergent application from the 4 to 6 true leaf stage of crop | |
| Tank Mix Components | AC 299,263 120 AS Herbicide Solution | 167 mL/ha |
| | Starane or Starane II Herbicide | 0.6 L/ha Starane OR 0.31 L/ha Starane II |
| | 2,4-D Ester | 560 g ae/ha (2,4-D Ester 600 formulation: 1 L/ha) (2,4-D Ester 700 formulation: 0.85 L/ha) |
| Adjuvant | Refer to table of Adjuvant Options above. Adjuvant option and rate will depend on wheat variety. | |
| Spray Volume | 50-100 L/ha water | |
| Weeds Controlled | <p>This tank-mix combination controls all weeds as listed on each of the tank mix partner product labels.</p> <p>Initial crop injury may be observed after application but this is outgrown and should not affect yield. Severe crop injury and a delay in maturity may occur as a result of over spray. AVOID SPRAYER OVERLAP.</p> | |

| AC 299,263 120 AS Herbicide Solution + Starane™ or Starane II Herbicide + CURTAIL® M Herbicide | | |
|---|--|--|
| Crops | Clearfield and Clearfield Plus Wheat ¹ | |
| Timing | Post-emergent application from the 3 to 6 true leaf stage of crop | |
| Tank Mix Components | AC 299,263 120 AS Herbicide Solution | 167 mL/ha |
| | Starane or Starane II Herbicide | 0.6 L/ha Starane OR 0.31 L/ha Starane II |
| | CURTAIL M | Maximum rate of 2.0 L/ha |
| Adjuvant | Refer to table of Adjuvant Options above. Adjuvant option and rate will depend on wheat variety. | |
| Spray Volume | 50-100 L/ha water | |
| Weeds Controlled | <p>This tank-mix combination controls all weeds as listed on each of the tank mix partner product labels.</p> <p>Initial crop injury may be observed after application but this is outgrown and should not affect yield. Severe crop injury and a delay in maturity may occur as a result of over spray. AVOID SPRAYER OVERLAP.</p> | |

| AC 299,263 120 AS Herbicide Solution + MCPA Ester | | |
|--|--|---|
| Crops | Clearfield and Clearfield Plus Wheat ¹ | |
| Timing | Post-emergent application from the 3 to 6 true leaf stage of crop | |
| Tank Mix Components | AC 299,263 120 AS Herbicide Solution | 167 mL/ha |
| | MCPA Ester | 560 g ai/ha – 630 g ai/ha (500 g/L formulation: 1.12 – 1.26 L/ha) (600 g/L formulation: 0.93 – 1.05 L/ha) |
| Adjuvant | Refer to table of Adjuvant Options above. Adjuvant option and rate will depend on wheat variety. | |
| Spray Volume | 50-100 L/ha water | |
| Weeds Controlled | <p>This tank-mix combination controls all weeds as listed on the AC 299,263 120 AS Herbicide Solution label plus weeds as listed on the MCPA Ester label.</p> <p>Initial crop injury may be observed after application but this is outgrown and should not affect yield. Severe crop injury and a delay in maturity may occur as a result of over spray. AVOID SPRAYER OVERLAP.</p> | |

CROP: Peas (Dried Field)

AC 299,263 120 AS Herbicide Solution + Basagran Forte Herbicide

AC 299,263 120 AS Herbicide Solution may be applied as an early post-emergence treatment in peas (dried field) when applied in tank mix combination with Basagran Forte Herbicide.

Follow all recommendations, precautions and restrictions on the tank-mix partner label.

Treated crops are not to be grazed or cut for hay.

| Timing | Early post-emergence | | | | | |
|------------------------------------|---|----------------|--------------------|------------------------------------|----------------|----------------------------------|
| Rate | 167 mL/ha AC 299,263 120 AS Herbicide Solution + 894 mL/ha Basagran Forte Herbicide + 2 L/ha nitrogen source* (UAN 28%) | | | | | |
| Water Volume | 100 L/ha | | | | | |
| Weeds Controlled | <p>This tank-mix combination will provide control of weeds as listed in the Broadleaf Weed and Grass Control section of this label. The addition of 894 mL/ha of Basagran Forte Herbicide will provide control of the following additional weeds:</p> <table border="1"><thead><tr><th>Control</th><th>Suppression</th></tr></thead><tbody><tr><td>Volunteer Clearfield canola</td><td rowspan="2">Wild buckwheat</td></tr><tr><td>Wild mustard (Group 2 resistant)</td></tr></tbody></table> | Control | Suppression | Volunteer Clearfield canola | Wild buckwheat | Wild mustard (Group 2 resistant) |
| Control | Suppression | | | | | |
| Volunteer Clearfield canola | Wild buckwheat | | | | | |
| Wild mustard (Group 2 resistant) | | | | | | |
| Remark | Application should be made from the 3 - 6 node stage of field peas and after weeds have emerged. Apply when broadleaf weeds are from the cotyledon to 4-leaf stage and when grassy weeds are at the 1 - 4 true leaf or early tillering. Initial transient crop yellowing may be observed after application but this is outgrown and should not affect yield. | | | | | |

*Nitrogen source: A reduction in grass control can be observed without the addition of a nitrogen source (UAN 28%).

MIXING INSTRUCTIONS

Use 50-100 L/ha of water. Use a 50-mesh (or coarser) filter screen. Fill the spray tank with three-quarters of the required amount of clean water, start agitation and continue agitation throughout the entire mixing and spraying procedure. Add the required amount of AC 299,263 120 AS Herbicide Solution into the sprayer through the tank opening and continue agitating. If required, add tank mix partner(s) and continue agitation. After the herbicide is thoroughly mixed, continue agitation and add the required amount of adjuvant or nitrogen source. If excess foaming occurs, a silicone anti-foaming agent may be added (e.g. HALT®). Complete filling the tank to the desired level with water. If agitation is stopped for more than 5 minutes, re-suspend spray solution by full agitation prior to commencing spraying again. Between loads of AC 299,263 120 AS Herbicide Solution, check in-line and nozzle screens and rinse and clean if necessary. Upon completion of

spraying AC 299,263 120 AS Herbicide Solution, thoroughly flush tank, boom, hoses and in-line and nozzle screens with clean water to avoid possible injury to other crops.

FOLLOW CROPPING

Winter wheat can be planted 3 months after treatment (3 MAT) as a rotational crop.*

Initial crop injury to non-**Clearfield** canola may be observed. Avoid spray overlap as yield reduction may result. The following crops may be grown safely the year following an application:

- Field peas
- Field corn
- Canary seed*
- Clearfield** Canola¹ (canola varieties with the **Clearfield** trait)
- Non-**Clearfield** canola*
- Lentils
- Spring wheat
- Durum wheat*
- Spring barley
- Sunflower
- Tame oats*
- Flax*
- Chickpea

The following crop may be grown safely two years following an application:

- Mustard (condiment type only)*

*If drought conditions are experienced between June 1 and September 1 in the year of application, delay planting of winter wheat, durum wheat, canary seed, tame oats, flax and canola (non-**Clearfield**) by an additional year. If drought is received between June 1 and September 1 in the year of application OR between June 1 and September 1 in the year following application, delay planting of mustard by an additional year.

In the grey, black, and dark brown soil zones, drought is defined as less than 125 mm of total precipitation between June 1 and September 1. In the brown soil zone, drought is defined as less than 125 mm of total precipitation between June 1 and September 1, OR less than 15 mm of precipitation in any month between June 1 and September 1.

There are insufficient data for other follow crops. Conduct a field bioassay (a test strip grown to maturity) the year before growing any crop other than those listed above.

WARNING: Certain environmental conditions may delay the breakdown of herbicide residues in soil. These conditions include but are not limited to drought, extremes in soil pH (less than 6.5), low organic matter soils (less than 3%), extremely coarse textured (sandy) soils, and excessive cold. Under these conditions, the level of phytotoxic herbicide residues present in the field the season following an application may result in an increased potential for injury to succeeding crops to occur. This potential for increased residues under these environmental conditions is not unique to any specific herbicide or herbicide group but is a property of those herbicides which persist in

the soil and are dependent on soil microbial activity and other non-microbial processes (e.g., hydrolysis) to breakdown.

RESISTANCE-MANAGEMENT RECOMMENDATIONS

For resistance management, AC 299,263 120 AS Herbicide Solution is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to AC 299,263 120 AS Herbicide Solution and other Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of AC 299,263 120 AS Herbicide Solution or other Group 2 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact BASF at 1-877-371-2273 or at www.agsolutions.ca.

® All other products listed are registered trademarks of their respective companies.

¹ *Methods of growing plants which are resistant to certain acetohydroxyacid synthase (AHAS) herbicides are protected by Canadian Patent No. 1341465. Other applicable patents include Canadian Patent Nos. 2456314, 2455512, 2456311, 2492167 and 2790951. The use of this product in the practice of patented methods could constitute patent infringement. Further information can be obtained by contacting BASF Canada at 1-877-371-2273.*

(Container)



Starane™ II

Herbicide

| | | |
|-------|---|-----------|
| GROUP | 4 | HERBICIDE |
|-------|---|-----------|

Starane II Herbicide is a selective herbicide for postemergent control of annual broadleaved weeds including cleavers and certain ALS-resistant broadleaved weeds such as kochia in spring wheat, durum wheat, winter wheat, spring barley, oats, rangeland, permanent pasture, rights-of-way, industrial and other non-crop areas of Canada. Starane II Herbicide will also suppress hemp-nettle, common chickweed and wild buckwheat.

COMMERCIAL (AGRICULTURAL)

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENT: fluroxypyr, present as 1-methylheptyl ester - 333 g/L
Emulsifiable concentrate

REGISTRATION NO. 29463

PEST CONTROL PRODUCTS ACT

WARNING
EYE AND SKIN IRRITANT
POTENTIAL SKIN SENSITIZER

NET CONTENTS: 1 L - bulk

Corteva Agriscience Canada Company
Suite 240, 115 Quarry Park Rd. SE
Calgary, Alberta
T2C 5G9
1-800-667-3852

™ Trademark of Corteva Agriscience and its affiliated companies.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN

Causes eye irritation. DO NOT get in eyes. May irritate the skin. Avoid contact with skin. Potential skin sensitizer.

For agricultural/crop scenarios: Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12-hours.

For non-crop areas: Do not enter or allow worker entry into treated areas until sprays have dried.

Do not use in residential areas. Residential areas are defined as any use site where bystanders including children could be exposed during or after application. This includes homes, schools, parks, playgrounds, playing fields, public buildings, or any other area where the general public including children could be exposed.

Apply only when potential for drift to areas of human habitation and human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversion, application equipment and sprayer settings.

PROTECTIVE CLOTHING AND EQUIPMENT

Wear a hat, coveralls over a long-sleeved shirt and long pants, socks, rubber boots or work boots and chemical-resistant gloves during spraying or application. In addition, wear goggles or face shield during mixing, loading, repair and clean-up or when handling the concentrate. Applicators using a closed cab are not required to wear chemical-resistant gloves.

OPERATOR USE PRECAUTIONS

Wear freshly laundered clothing and clean protective equipment daily. At completion of spraying or end of the day take a shower immediately. Wash thoroughly with soap and water before eating, drinking, smoking or using the toilet. Remove contaminated clothing at the end of the work session and store and wash separately from household laundry using detergents and hot water before reuse. If herbicide penetrates clothing, remove immediately; then wash thoroughly and put on clean clothing. Throw away clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Avoid contact with eyes, skin and clothing. Avoid breathing spray mist.

PHYSICAL OR CHEMICAL HAZARDS

COMBUSTIBLE. Do not use or store near heat or open flame.

FIRST AID

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION

THIS PRODUCT CONTAINS A PETROLEUM DISTILLATE. Vomiting may cause aspiration pneumonia. The decision of whether to induce vomiting or not should be made by the attending physician. If lavage is performed, suggest endotracheal and/or oesophageal control. Danger from lung aspiration of petroleum based solvents must be weighed against toxicity when considering emptying the stomach. No specific antidote. Employ supportive care. Treatment should be based on the judgment of the physician in response to reactions of the patient.

AGRICULTURAL CHEMICAL

Do not ship or store with food, feeds, drugs or clothing.

ENVIRONMENTAL PRECAUTIONS

TOXIC to non-target terrestrial plants and aquatic organisms. Observe spray buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

This product contains an active ingredient and aromatic petroleum distillates which are toxic to aquatic organisms.

STORAGE

Store in original containers in a secure, dry heated storage. If product is frozen, bring to room temperature and agitate before use. Do not allow contamination of seeds, plants, fertilizers or other pesticides. To prevent contamination, store this product away from food or feed. If containers are damaged or spill occurs, use the product immediately or contain the spill with absorbent materials and dispose of waste.

DISPOSAL**Recyclable Containers**

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/territorial requirements.

Returnable Containers

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

Refillable Containers

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

The user assumes the risk to persons or property that arises from any such use of this product.

(Booklet)



Starane™ II

Herbicide

| | | |
|-------|---|-----------|
| GROUP | 4 | HERBICIDE |
|-------|---|-----------|

Starane II Herbicide is a selective herbicide for postemergent control of annual broadleaved weeds including cleavers and certain ALS-resistant broadleaved weeds such as kochia in spring wheat, durum wheat, winter wheat, spring barley, oats, rangeland, permanent pasture, rights-of-way, industrial and other non-crop areas of Canada. Starane II Herbicide will also suppress hemp-nettle, common chickweed and wild buckwheat.

COMMERCIAL (AGRICULTURAL)

READ THE LABEL AND BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENT: fluroxypyr, present as 1-methylheptyl ester - 333 g/L
Emulsifiable concentrate

REGISTRATION NO. 29463 PEST CONTROL PRODUCTS ACT

WARNING
EYE AND SKIN IRRITANT
POTENTIAL SKIN SENSITIZER

NET CONTENTS: 1 L - bulk

Corteva Agriscience Canada Company
Suite 240, 115 Quarry Park Rd. SE
Calgary, Alberta
T2C 5G9
1-800-667-3852

™Trademark of Corteva Agriscience and its affiliated companies.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN

Causes eye irritation. DO NOT get in eyes. May irritate the skin. Avoid contact with skin. Potential skin sensitizer.

For agricultural/crop scenarios: Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12-hours.

For non-crop areas: Do not enter or allow worker entry into treated areas until sprays have dried.

Do not use in residential areas. Residential areas are defined as any use site where bystanders including children could be exposed during or after application. This includes homes, schools, parks, playgrounds, playing fields, public buildings, or any other area where the general public including children could be exposed.

Apply only when potential for drift to areas of human habitation and human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversion, application equipment and sprayer settings.

PROTECTIVE CLOTHING AND EQUIPMENT

Wear a hat, coveralls over a long-sleeved shirt and long pants, socks, rubber boots or work boots and chemical-resistant gloves during spraying or application. In addition, wear goggles or face shield during mixing, loading, repair and clean-up or when handling the concentrate. Applicators using a closed cab are not required to wear chemical-resistant gloves.

OPERATOR USE PRECAUTIONS

Wear freshly laundered clothing and clean protective equipment daily. At completion of spraying or end of the day take a shower immediately. Wash thoroughly with soap and water before eating, drinking, smoking or using the toilet. Remove contaminated clothing at the end of the work session and store and wash separately from household laundry using detergents and hot water before reuse. If herbicide penetrates clothing, remove immediately; then wash thoroughly and put on clean clothing. Throw away clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Avoid contact with eyes, skin and clothing. Avoid breathing spray mist.

PHYSICAL OR CHEMICAL HAZARDS

COMBUSTIBLE. Do not use or store near heat or open flame.

FIRST AID

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION

THIS PRODUCT CONTAINS A PETROLEUM DISTILLATE. Vomiting may cause aspiration pneumonia. The decision of whether to induce vomiting or not should be made by the attending physician. If lavage is performed, suggest endotracheal and/or oesophageal control. Danger from lung aspiration of petroleum based solvents must be weighed against toxicity when considering emptying the stomach. No specific antidote. Employ supportive care. Treatment should be based on the judgment of the physician in response to reactions of the patient.

AGRICULTURAL CHEMICAL

Do not ship or store with food, feeds, drugs or clothing.

ENVIRONMENTAL PRECAUTIONS

TOXIC to non-target terrestrial plants and aquatic organisms. Observe spray buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

This product contains an active ingredient and aromatic petroleum distillates which are toxic to aquatic organisms.

STORAGE

Store in original containers in a secure, dry heated storage. If product is frozen, bring to room temperature and agitate before use. Do not allow contamination of seeds, plants, fertilizers or other pesticides. To prevent contamination, store this product away from food or feed. If containers are damaged or spill occurs, use the product immediately or contain the spill with absorbent materials and dispose of waste.

DISPOSAL**Recyclable Containers**

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/territorial requirements.

Returnable Containers

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

Refillable Containers

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean-up of spills.

GENERAL INFORMATION

Starane II Herbicide is a selective postemergent herbicide for the control of annual broadleaved weeds in spring wheat, durum wheat, winter wheat, spring barley and oats, not underseeded with legumes. Starane II Herbicide offers a novel mode of action for controlling hard-to-kill annual broadleaved weeds such as kochia (including ALS resistant biotypes) and cleavers. Starane II Herbicide will also provide suppression of hempnettle, wild buckwheat and chickweed.

MODE OF ACTION

Starane II Herbicide is a systemic auxin-type herbicide and moves within the plant for control of exposed and underground plant tissues. The product controls weeds by disrupting normal plant growth patterns. Symptoms of weeds include epinasty (twisting of the stems) and swollen nodes.

GENERAL USE PRECAUTIONS**Sensitive Plants**

Do not apply Starane II Herbicide directly to, or otherwise permit it to come in direct contact with susceptible crops or desirable plants including alfalfa, edible beans, canola, flowers and ornamentals, lentils, lettuce, peas, potatoes, radishes, soybeans, sugar beets, sunflowers, tomatoes or tobacco.

Non-Target Sites

Do not apply where proximity of susceptible crops or other desirable plants is likely to result in exposure to spray or spray drift. See Environmental Precautions section of the label.

Crop Rotation

Fields previously treated with Starane II Herbicide can be seeded the following year to alfalfa, barley, canola, corn, dry beans, fababeans, flax, forage grasses, lentils, mustard, oats, peas, potatoes, rye, soybeans, sugar beets, sunflowers or wheat or fields can be summerfallowed.

Tank Mixtures

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Corteva Agriscience Canada Company at 1-800-667-3852 or www.corteva.ca for information before applying any tank mix that is not specifically recommended on this label.

Spray Equipment Precaution

Do not apply through any type of irrigation system.

To Reduce Spray Drift

- Use nozzles delivering higher volumes and coarser droplets.
- Use low pressures (200 to 275 kPa).
- Spray when the wind velocity is 15 km/hr or less.
- Spot treatments should only be applied with a calibrated boom to prevent over-application.

Equipment Clean-Up

Equipment used to apply Starane II Herbicide should not be used to apply other pesticides to sensitive crops without thorough cleaning. Contact your Starane II Herbicide dealer for detailed equipment cleaning procedures.

DIRECTIONS FOR USE

READ THE ENTIRE LABEL BEFORE USE. FAILURE TO FOLLOW LABEL INSTRUCTIONS MAY RESULT IN ERRATIC WEED CONTROL OR CROP DAMAGE. DO NOT APPLY TO CROPS UNDERSEEDING WITH LEGUMES.

As this product is not registered for the control of pests in aquatic systems, **DO NOT** use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Field sprayer application

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Boom height must be 60 cm or less above the crop or ground.

Aerial application

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply when wind speed is greater than 16 km/h at flying height at the site of application. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Reduce drift caused by turbulent wingtip vortices. Nozzle distribution along the spray boom length MUST NOT exceed 65% of the wing or rotorspan.

Crops Registered

spring wheat
durum wheat
winter wheat
spring barley
oats

STARANE II HERBICIDE ALONE

Apply Starane II Herbicide with ground or aerial application equipment (see Application directions section) for postemergent control of cleavers and kochia, including ALS resistant kochia and other broadleaved weeds infesting spring wheat, durum wheat, winter wheat, spring barley and oats. See following table for weeds controlled/suppressed and applications rates.

Weeds Controlled Or Suppressed By Starane II Herbicide Alone

| Weeds Controlled | Starane II Herbicide Rate (L/ha) |
|--------------------------------|----------------------------------|
| chickweed* (up to 8 cm) | 0.41 |
| cleavers (1-6 whorls) | 0.21 |
| cleavers (1-8 whorls) | 0.31 |
| kochia* (2-8 leaf) | 0.31 |
| round-leaved mallow (1-6 leaf) | 0.41 |
| volunteer flax (1-12 cm) | 0.31 |
| Weeds Suppressed | Starane II Herbicide Rate (L/ha) |
| hempnettle (2-6 leaf) | 0.41 |
| stork's-bill (1-8 leaf) | 0.31 |
| wild buckwheat (1-8 leaf) | 0.21 |

*Including biotypes resistant to Group 2 herbicides that inhibit the ALS enzyme.

NOTE: Starane II Herbicide activity is influenced by weather conditions. Optimum activity requires active crop and weed growth. The temperature range for optimum activity is 12°C to 24°C. Reduced activity will occur when temperatures are below 8°C or above 27°C. Frost before application (3 days) or shortly after (3 days) may reduce weed control and crop tolerance. Weed control may be reduced during stress

conditions, e.g. drought, heat or cold stress, or if weeds have initiated flowering, or if heavy infestations exist.

Application Timing

Apply to actively growing spring wheat, durum wheat, spring barley and oats from the 2-leaf crop growth stage up to and including initiation of stem elongation. Apply to winter wheat in the spring from the 3 tiller stage to just before the flag leaf stage. Apply when weeds are actively growing. Best results are obtained from applications made to seedling weeds. Only weeds emerged at the time of treatment will be controlled. Extreme growing conditions such as drought or near freezing temperature prior to, at and following time of application may reduce weed control and increase the risk of crop injury at all stages of growth. Foliage that is wet at the time of application may decrease control. Starane II Herbicide applications are rainfast within 1 hour after application. Under conditions of low crop and high weed density, control may be reduced.

Preharvest/Grazing Intervals (Starane II Herbicide applied alone)

- Livestock may be grazed on treated crops 3 days following application.
- Do not harvest the treated crop within 60 days after application.

APPLICATION DIRECTIONS (GROUND AND AERIAL APPLICATIONS)

(1) Ground Application

Using ground equipment, apply Starane II Herbicide alone or with tank mix partners as a broadcast treatment. Apply Starane II Herbicide alone at the recommended rates in a spray volume of 50-100 L water/ha.

(2) Aerial Application

Do not use human flaggers.

Use Starane II Herbicide alone or tank mixed with a product registered for aerial application as a broadcast treatment. Apply the recommended rates of Starane II Herbicide and tank mix partner as indicated elsewhere on this label. Apply in a spray volume of 30-50 L water/ha.

Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate swath marking devices.

Use Precautions

Apply only when meteorological conditions at the treatment site allow for complete and even coverage. Apply only under conditions of good practice specific to aerial application as outlined in the National Aerial Pesticide Application Manual, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

Operator Precautions

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

Product Specific Precautions

Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-800-667-3852 or obtain technical advice from the distributor or your provincial agricultural representative. Application of this specific product must meet and/or conform to the following:

- The interaction of many equipment-and-weather-related factors determine the potential for spray drift. Users are responsible for considering all these factors when making decisions.

The following drift management requirement must be followed to avoid off-target drift movement from aerial applications: Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

TANK-MIX COMBINATIONS - STARANE II HERBICIDE + LIQUID ACHIEVE¹ SC HERBICIDE OR ACHIEVE LIQUID HERBICIDE (Do not apply to oats)

For control of wild oats and green foxtail in spring wheat, winter wheat and spring barley apply 0.5 L/ha of Liquid Achieve SC or Achieve Liquid with Starane II Herbicide. Refer to the above table for rates and broadleaved weeds controlled or suppressed with Starane II Herbicide. To control additional broadleaved weeds tank mix with either 2,4-D ester or MCPA ester up to a maximum of 560 g ae/ha. (equivalent to 1.0 L/ha of 2,4-D LV600 Herbicide (0.85 L/ha 2,4-D Ester 700 Herbicide) or 1.12 L/ha of MCPA LV500 Herbicide) (0.93 L/ha MCPA Ester 600 Herbicide) or 1.5 to 2.0 L/ha Curtail M Herbicide. Refer to the 2,4-D ester or MCPA ester tank mix partner label for rates less than 560 g ae/ha and weeds controlled.

Mixing Instructions

1. Begin to fill sprayer tank with clean water, and engage agitator. **Agitation must be continued throughout the entire mixing and spraying procedure.**
2. When the sprayer is three quarters full of water, add Liquid Achieve SC or Achieve Liquid. If more than one container of Liquid Achieve SC or Achieve Liquid is to be used, add all Liquid Achieve SC or Achieve Liquid containers prior to adding Starane II Herbicide.
3. Add Starane II Herbicide next. If more than one container of Starane II Herbicide is to be used, add all Starane II Herbicide containers prior to adding any other tank-mix partners.
4. If 2,4-D ester, MCPA ester or Curtail M are also being added, add these next. If more than one container of 2,4-D ester, MCPA ester or Curtail M is to be used, add all 2,4-D ester, MCPA ester or Curtail M containers prior to adding Turbocharge D or Turbocharge.
5. Lastly, add the required amount of Turbocharge D or Turbocharge and then continue to fill tank to desired level with water.

TANK-MIX COMBINATIONS - STARANE II HERBICIDE + CURTAILTM M HERBICIDE

Starane II Herbicide mixed with Curtail M Herbicide will control a wide spectrum of broadleaf weeds.

Mixing Instructions

Add the correct amount of Starane II Herbicide to spray tank 1/2 filled with water and agitate. Add the correct amount of Curtail M Herbicide and continue to agitate.

Timing of Application

Apply postemergence when weeds are in the seedling stage (2-4 leaf) and when spring wheat, durum wheat, spring barley and oats are in the 3-leaf to just before the flag leaf emergence stage. Apply to

winter wheat in the spring from the 3 tiller stage to just before the flag leaf stage. Do not apply later than the flag-leaf stage.

Low Rate: To control or suppress the following weeds apply a uniform spray containing Starane II Herbicide at 0.31 L/ha tank mixed with Curtail M Herbicide at 1.5 L/ha.

Weeds Controlled

| | |
|-------------------------------------|---------------------------|
| Canada thistle (low infestations) | mustard, wild |
| cleavers (1-8 whorls) | shepherd's-purse |
| flax, volunteer (1-12 cm) | stinkweed |
| flixweed (spring seedling 2-4 leaf) | sunflower, volunteer |
| kochia*** | wild buckwheat (1-8 leaf) |
| lamb's-quarters | |

Weeds Suppressed

stork's-bill (1-8 leaf)

High Rate: To control or suppress the following weeds apply a uniform spray containing Starane II Herbicide at 0.41 L/ha tank mixed with Curtail M Herbicide at 2.0 L/ha.

Weeds Controlled

| | | |
|------------------------------|--------------------------------|---------------------------|
| Canada thistle* | lamb's-quarters | sow-thistle (perennial) * |
| canola (volunteer) | redroot pigweed | stinkweed |
| cleavers (1-8 whorls) | round-leaved mallow (1-6 leaf) | stork's-bill (1-8 leaf) |
| common chickweed* | Russian pigweed | sunflower (volunteer) |
| common groundsel | scentless chamomile | tartary buckwheat |
| dandelion** | shepherd's-purse | volunteer flax (1-12 cm) |
| flixweed** | smartweed | wild buckwheat (1-8 leaf) |
| hemp-nettle (2-6 leaf stage) | sow-thistle (annual) | wild mustard |
| kochia*** | | |

* Season long control, with some regrowth in the fall (top growth control).

** Spring rosettes only.

*** Including biotypes resistant to Group 2 herbicides that inhibit the ALS enzyme.

The active ingredients in this tank mixture will only control chickweed that is emerged at TIME OF application. Chickweed plants which emerge after application will not be controlled. To improve the reduction in chickweed population at the end of season, delay the timing of application as late as possible to when the majority of chickweed plants have emerged. The reduction in chickweed population will also be improved in crops which are more competitive and which allow limited light penetration.

Pre-Harvest/Grazing Intervals (Starane II Herbicide + Curtail M Herbicide)

- Do not cut or graze treated fields within 7 days after application.
- Do not harvest the treated mature crop within 60 days after application.

TANK MIXES WITH STARANE II HERBICIDE + CURTAIL M HERBICIDE FOR ANNUAL GRASS CONTROL (Do not apply to oats)

Tank mixtures of Starane II Herbicide + Curtail M Herbicide with other herbicides will provide control of additional weeds. See following table for tank mixes with Starane II Herbicide + Curtail M Herbicide.

Tank-Mix Combinations with Starane II Herbicide at 0.31 L/ha + Curtail M Herbicide at 1.5 L/ha or Starane II Herbicide at 0.41 L/ha + Curtail M Herbicide at 2.0 L/ha

| Herbicide Tank-Mix Partner | Crops Registered | Rate of Tank-Mix Partner | Adjuvant Rate | Additional Weeds |
|-------------------------------------|--|--------------------------|--|----------------------------|
| Liquid Achieve SC or Achieve Liquid | spring wheat, durum wheat, winter wheat, spring barley | 0.5 L/ha | Turbocharge D or Turbocharge 0.5 L/100 L of spray volume (0.5% v/v) | green foxtail wild oats |

TANK-MIX COMBINATION - STARANE II HERBICIDE + 2,4-D ESTER HERBICIDE (Do not apply to oats)

For control of a wide spectrum of broadleaved weeds (see following table) apply Starane II Herbicide tank mixed with 2,4-D ester herbicide. Apply postemergence when weeds are in the seedling stage (2-4 leaf) and when spring wheat, durum wheat, winter wheat and spring barley are in the 4-leaf to flag-leaf stage. Apply to winter wheat in the spring from the 3 tiller stage to just before the flag leaf stage. Application before the 4-leaf stage of wheat and barley may cause severe twisting of leaves and stem, and head deformities which may reduce yield up to 10%. Do not apply later than the flag-leaf stage.

Rates of Applications

I. Low Rate: To control the following weeds apply a uniform spray containing 0.23 L/ha Starane II Herbicide and 420 g a.e./ha 2,4-D ester herbicide (equivalent to 0.64 L/ha of 2,4-D Ester LV700 Herbicide).

Weeds Controlled

| | | |
|-----------------------|--------------------------------------|---------------------------|
| bluebur | goat's-beard | shepherd's-purse |
| burdock | hoary cress* | stinkweed |
| canola (volunteer) | kochia*** | sunflower (annual) |
| cleavers (1-6 whorls) | lamb's-quarters | vetch |
| clovers (sweet) | mustards (except green, dog & tansy) | wild radish |
| cocklebur | plantain | wild mustard |
| flixweed | prickly lettuce | wild buckwheat (1-6 leaf) |
| field horsetail* | ragweed | |

II. High Rate: To control the above weeds plus the additional weeds listed below, apply a uniform spray containing 0.31 L/ha of Starane II Herbicide and 560 g a.e./ha of 2,4-D ester herbicide (equivalent to 0.85 L/ha of 2,4-D Ester LV700 Herbicide).

Additional Weeds Controlled:

| | | |
|-----------------------|--------------------------------|---------------------------|
| blue lettuce* | hairy galinsoga | Russian thistle |
| cleavers (1-8 whorls) | hedge bindweed | smartweed |
| dandelion** | lady's-thumb | stork's-bill (1-8 leaf) |
| docks | leafy spurge* | tansy mustard |
| dog mustard | oak-leaved goosefoot | tartary buckwheat |
| field bindweed* | redroot pigweed | volunteer flax (1-12 cm) |
| field peppergrass | round-leaved mallow (1-6 Leaf) | wild buckwheat (1-8 leaf) |
| gumweed | | |

Weeds Suppressed:

common chickweed*** (up to 8 cm) hemp-nettle (2-6 leaf stage) sow-thistle (perennial) ♦
Canada thistle♦ sow-thistle (annual)

♦ Top growth control only

♦♦ Spring rosettes

♦♦♦ Including biotypes resistant to Group 2 herbicides that inhibit the ALS enzyme

NOTE: The activity of Starane II Herbicide combined with 2,4-D ester herbicide is influenced by weather conditions. Optimum activity requires active crop and weed growth. The temperature range for optimum activity is 12°C to 24°C. Reduced activity will occur when temperatures are below 8°C or above 27°C. Frost before application (3 days) or shortly after (3 days) may reduce weed control and crop tolerance. Weed control may be reduced during stress conditions, e.g. drought, heat or cold stress, or if weeds have initiated flowering, or if heavy infestations exist.

Mixing Instructions

Add the correct amount of Starane II Herbicide to spray tank 1/2 filled with water and agitate. Add the correct amount of 2,4-D ester herbicide (e.g. 2,4-D Ester LV700 Herbicide) and continue to agitate.

Pre-Harvest/Grazing Intervals (Starane II Herbicide + 2,4-D ester herbicide)

- Do not permit lactating dairy animals to graze fields within 7 days after application.
- Do not harvest forage or cut hay within 30 days after application.
- Withdraw meat animals from treated fields at least 3 days before slaughter.
- Do not harvest the treated mature crop within 60 days after application.

TANK-MIX COMBINATIONS - STARANE II HERBICIDE + 2,4-D ESTER HERBICIDE + OTHER HERBICIDES (Do not apply to oats)

Tank mixtures of Starane II Herbicide + 2,4-D ester herbicide with other herbicides will provide control of additional weeds. For products which may be tank mixed with Starane II Herbicide + 2,4-D ester herbicide and the additional weeds controlled see following table.

Tank-Mix Combinations with Starane II Herbicide (0.31 L/ha) + 2,4-D ester herbicide (560 g a.e./ha) or Starane II Herbicide (0.23 L/ha) + 2,4-D ester herbicide (420 g a.e./ha)

| Herbicide Tank-Mix Partner | Crops Registered | Rate of Tank-Mix Partner | Adjuvant Rate | Additional Weeds |
|--|--|--------------------------|---|---|
| Liquid Achieve SC or Achieve Liquid*♦♦ | spring wheat, durum wheat, winter wheat, spring barley | 0.5 L/ha | Turbocharge D or Turbocharge 0.5 L/100 L of spray volume (0.5% v/v) | green foxtail, wild oats |
| Simplicity™♦♦ | spring wheat, durum wheat, winter wheat | 0.5 L/ha | None required | barnyard grass (1-5 leaf), common chickweed (up to 10 cm), hemp nettle (1-8 leaf), Japanese brome (1-6 leaf), wild oats (up to 4 leaf, 2 tiller), wild buckwheat (1-4 leaf) |

*This tank-mix may cause temporary crop injury. However, yield will not normally be affected. This tank-mix may reduce control of wild oats compared to Liquid Achieve SC or Achieve Liquid Herbicide alone.

♦♦Tank mixing with Simplicity only applies to the low rate of Starane II Herbicide + 2,4-D ester

TANK-MIX COMBINATIONS – STARANE II HERBICIDE + DRAFT OR REFINE SG HERBICIDE OR REFINE SG TOSS-N-GO HERBICIDE

For control of a wide spectrum of broadleaved weeds (see following table) Starane II Herbicide can be tank mixed with Draft or Refine SG Herbicide or Refine SG Toss-N-Go Herbicide. Apply post-emergence when weeds are actively growing, with best results on seedling weeds that are less than 10 cm in height. Only weeds emerged at the time of treatment will be controlled. Thorough covering of the target weeds is essential. The tank-mix can be applied to actively growing spring wheat, including durum, spring barley and oats from the 2-leaf crop growth stage up to and including initiation of stem elongation. The tank-mix of Draft or Refine SG Herbicide or Refine SG Toss-N-Go Herbicide with Starane II Herbicide allows for the control of Group 2 resistant cleavers at 0.21 L/ha and kochia 0.31 L/ha. Do not apply to crops underseeded to legumes or grasses.

Rates of Application To control the following weeds apply a uniform spray containing 0.21 to 0.41 litres of Starane II Herbicide with 20 g/ha Draft or 30 g/ha of Refine SG Herbicide or Refine SG Toss-N-Go Herbicide and 2 litres per 1000 L spray solution of non-ionic surfactant (Agral 90, Ag-surf or Citowett Plus) with a spray volume of 100 L/ha.

Weeds Controlled or Suppressed

Weeds Controlled at 0.21 L/ha of Starane II Herbicide and 20 g/ha Draft or 30 g/ha of Refine SG Herbicide or Refine SG Toss-N-Go Herbicide:

| | | |
|--|----------------------------|---------------------------------|
| annual smartweed (green smartweed, lady's thumb) | groundsel (common) | shepherd's purse |
| ball mustard | hemp-nettle | stinkweed |
| chickweed (1-6 leaf) ¹ | kochia ² | tartary buckwheat |
| cleavers (1-4 whorls) | lamb's-quarters | volunteer rapeseed ³ |
| corn spurry | narrow-leaved hawk's beard | volunteer sunflower |
| cow cockle | redroot pigweed | wild buckwheat ⁴ |
| flixweed | Russian thistle | wild mustard |

¹ Spray before canopy prevents thorough coverage of weeds. Chickweed emerging after application will not be controlled.

² Excluding Group 2 resistant kochia

³ The tank-mix will not control imazethapyr tolerant canola varieties (e.g. Clearfield varieties or other varieties with the Pursuit Smart trait)

⁴ Apply to actively growing wild buckwheat at the cotyledon to 3-leaf stage. Control may be reduced under environmental stress. Large plants may regrow after treatment.

Weeds suppressed at 0.21 L/ha Starane II Herbicide and 20 g/ha Draft or 30 g/ha of Refine SG Herbicide or Refine SG Toss-N-Go Herbicide:

| | |
|---|-------------------------------------|
| Canada thistle ⁵ | sow thistle ⁵ |
| round leaved mallow (2-6 leaf) ⁶ | stork's bill (2-6 leaf stage) |
| scentless chamomile | toadflax (up to 15 cm) ⁷ |

⁵ Apply when most of the thistles have emerged and are actively growing. Apply before bud stage and before crop canopy prevents thorough coverage of thistles, when thistles are no larger than 15 cm in height for best top growth control. A single application will effectively inhibit the ability of emerged thistles to compete with the crop. Later emerging thistles will not be controlled.

⁶ Apply to actively growing round leaved mallow. Will keep mallow stunted, but may not be able to reduce overall populations

⁷ A control program for this weed includes chemical application and frequent tillage.

Additional Weeds Controlled at 0.31 L/ha of Starane II Herbicide and 20 g/ha Draft or 30 g/ha of Refine SG Herbicide or Refine SG Toss-N-Go Herbicide:

kochia (2-8 leaf; including group 2 resistant biotypes)
volunteer flax (1-12 cm)

Additional Weeds Controlled and Suppressed at 0.41 L/ha Starane II Herbicide and 20 g/ha Draft or 30 g/ha of Refine SG Herbicide or Refine SG Toss-N-Go Herbicide:

Weeds Controlled

round leaved mallow (1-6 leaf)

Weeds Suppressed

chickweed (up to 8 cm; including group 2 resistant biotypes)

Mixing Instructions

Fill the tank to 1/3 full of clean water, and with the agitator running, add the required amount of Draft or Refine SG Herbicide or Refine SG Toss-N-Go Herbicide and continue to agitate for a minimum of 5 minutes to ensure that Draft or Refine SG Herbicide or Refine SG Toss-N-Go Herbicide is completely dissolved. Add the required amount of Starane II Herbicide at a reduced agitation. Add the rest of the water and then add adjuvant. Use a spray volume of 100 L/ha.

Pre-Harvest/Grazing Intervals

- Livestock may not be grazed or fed crops within 7 days of application
- Do not harvest the treated crop within 60 days of application

TANK-MIX COMBINATIONS – STARANE II HERBICIDE + 2,4-D ESTER HERBICIDE + DRAFT OR REFINE SG HERBICIDE OR REFINE SG TOSS-N-GO HERBICIDE OR RETAIN A HERBICIDE (Do not apply to oats)

Spring Wheat (excluding durum) and spring barley

For control of a wide spectrum of broadleaf weeds, including cleavers, Starane II Herbicide may be tank mixed with 2,4-D ester herbicide and Draft or Refine SG Herbicide or Refine SG Toss-N-Go Herbicide or Retain A Herbicide. Apply Starane II Herbicide at 0.16 L/ha tank mixed with Draft at 20 g/ha, or Refine SG Herbicide or Refine SG Toss-N-Go Herbicide or Retain A Herbicide at 30 g/ha, 2,4-D ester herbicide at 280 g a.e./ha (equal to 0.5 L/ha of 2,4-D Ester LV 600 or 0.42 L/ha of 2,4-D Ester LV 700) and a non-ionic surfactant (Agral 90, Ag-Surf or Citowett Plus) at 2 L per 1000 L of spray solution. A spray volume of 100 L/ha should be utilized. Apply postemergence when weeds are in the seedling up to 4-leaf/whorls stage, and when spring wheat (excluding durum) and spring barley are in the 4-leaf to flag-leaf stage. **Do not apply to crops underseeded to legumes.**

Weeds Controlled or Suppressed

Weeds Controlled:

| | | |
|--|----------------------------|---------------------------------|
| annual smartweed (green smartweed, lady's thumb) | groundsel (common) | shepherd's purse |
| ball mustard | hemp-nettle | stinkweed |
| chickweed (1-6 leaf) ¹ | kochia ³ | tartary buckwheat |
| cleavers ² | lamb'-quarters | volunteer rapeseed ⁴ |
| corn spurry | narrow-leaved hawk's beard | volunteer sunflower |
| cow cockle | redroot pigweed | wild buckwheat ⁵ |
| flixweed | Russian thistle | wild mustard |

¹ Spray before crop canopy prevents thorough coverage of weeds. Chickweed emerging after application will not be controlled.

² Excluding Group 2 resistant cleavers

³ Excluding Group 2 resistant kochia

⁴ The tank-mix will not control imazethapyr tolerant canola varieties (e.g. Clearfield varieties or other varieties with the Pursuit Smart trait)

⁵ Apply to actively growing wild buckwheat from the cotyledon to 3-leaf stage. Control may be reduced under environmental stress. Large plants may regrow after treatment

Weeds Suppressed:

| | |
|---|--------------------------|
| Canada thistle ⁶ | sow thistle ⁶ |
| round leaved mallow (2-6 leaf) ⁷ | stork's bill (2-6 leaf) |

scentless chamomile

toadflax (to 15 cm)⁸

⁶ Apply when most of the thistles have emerged and are actively growing. Apply before bud stage, and before crop canopy prevents thorough coverage of thistles, when thistles are no larger than 15 cm in height for best top growth control. A single application will effectively inhibit the ability of emerged thistles to compete with the crop. Later emerging thistles will not be controlled.

⁷ Apply to actively growing round leaved mallow. Will keep mallow stunted, but may not be able to reduce overall populations.

⁸ A control program for this weed includes chemical application and frequent tillage.

Mixing Instructions

Fill the tank to 1/3 full of clean water. With agitator running, add required amount of Draft or Refine SG Herbicide or Refine SG Toss-N-Go Herbicide or Retain A Herbicide and continue to agitate for a minimum of 5 minutes to ensure the Draft or Refine SG Herbicide or Refine SG Toss-N-Go Herbicide or Retain A Herbicide is completely dissolved. After it is fully dissolved add the Starane II Herbicide with continuous agitation. Add the 2,4-D ester herbicide and the rest of the water. Then add adjuvant. Use a spray volume of 100 L/ha. Consult the 2,4-D label for additional application instructions and use precautions.

Preharvest/Grazing Intervals (Starane II Herbicide + Refine Extra or Draft + 2,4-D ester)

- Allow 7 days between application and grazing of lactating dairy animals.
- Do not harvest forage or cut hay within 30 days after application.
- Remove meat animals from treated fields at least 3 days prior to slaughter.
- Do not harvest the treated mature crop within 60 days after application.

TANK-MIX COMBINATION - STARANE II HERBICIDE + MCPA ESTER HERBICIDE

For control of a wide spectrum of broadleaved weeds (see following table) apply Starane II Herbicide at 0.31 L/ha tank mixed with MCPA LV500 Herbicide at 1.12 L/ha or MCPA Ester 600 Herbicide at 0.93 L/ha. Apply postemergence when weeds are in the seedling stage (2-4 leaf) and when spring wheat, durum wheat, spring barley and oats are in the 3-leaf to flag-leaf stage. Apply to winter wheat in the spring from the 3 tiller stage to just before the flag leaf stage. Application before the 3-leaf stage of wheat and barley may cause severe twisting of leaves and stem, and head deformities which may reduce yield up to 10%. Do not apply later than the flag-leaf stage. Do not apply to cereals underseeded with a legume.

Mixing Instructions

Add the correct amount of Starane II Herbicide to spray tank 1/2 filled with water and agitate. Add the correct amount of MCPA ester herbicide and continue to agitate.

Weeds Controlled or Suppressed by Starane II Herbicide+ MCPA ester herbicide Tank-Mix

Weeds Controlled

| | | |
|---------------------------|------------------------------------|---------------------------|
| burdock, common | kochia• | shepherd's-purse |
| canola, volunteer | lamb's-quarters | stinkweed |
| cleavers (1-8 whorls) | mustard (except dog & green tansy) | sunflower, annual |
| cocklebur | pigweed, redroot (1-4 leaf) | vetch |
| flax, volunteer (1-12 cm) | prickly lettuce | wild radish |
| flixweed | ragweed, common | wild buckwheat (1-8 leaf) |
| hempnettle (2-6 leaf) | | |

Weeds Suppressed

green smartweed
stork's-bill (1-8 leaf)

♦Including biotypes resistant to Group 2 herbicides that inhibit the ALS enzyme.

Pre-Harvest/Grazing Intervals (Starane II Herbicide + MCPA ester herbicide)

- Allow 7 days between application and grazing/cutting for hay.
- Do not harvest the treated mature crop within 60 days after application.

TANK-MIX COMBINATIONS - STARANE II HERBICIDE + MCPA ESTER HERBICIDE + OTHER HERBICIDES (Do not apply to oats)

Tank mixtures of Starane II Herbicide+ MCPA ester herbicide with other herbicides will provide control of additional weeds. For products which may be tank mixed with Starane II Herbicide + MCPA ester herbicide and the additional weeds controlled see following table.

Tank-Mix Combinations with Starane II Herbicide (0.31 L/ha) + MCPA LV500 Herbicide (1.12 L/ha) or MCPA Ester 600 Herbicide (0.93 L/ha)

| Herbicide Tank-Mix Partner | Crops Registered | Rate/ha | Adjuvant Rate | Additional Weeds |
|--------------------------------------|--|----------------|--|----------------------------|
| Liquid Achieve SC or Achieve Liquid* | spring wheat, durum wheat, winter wheat, spring barley | 0.5 L/ha | Turbocharge D or Turbocharge 0.5 L/ha (0.5% v/v) | green foxtail wild oats |

*MCPA ester can be tank mixed at rates up to a maximum of 560 g ae/ha with this tank-mix combination.

TANK-MIX COMBINATION - STARANE II HERBICIDE + SIMPLICITY™ (Do not apply to oats)

For control of a wide spectrum of broadleaf and grass weeds (see following table) in spring wheat, durum wheat and winter wheat, apply Starane II Herbicide tank mixed with Simplicity. Apply post-emergence when weeds are actively growing: grass weeds up to the 4 leaf, 2 tiller stage and broadleaf weeds as indicated in the following table. Best results are obtained from applications made to seedling weeds. Apply to actively growing spring or durum wheat from the 3 leaf stage to the first node stage. Apply to winter wheat in the 2-7 leaf, 4 tiller stage with a spring application. Extreme growing conditions such as drought or near freezing temperature prior to, at, or following time of application may reduce weed control and increase the risk of crop injury at all stages of growth. Under conditions of low crop and high weed density, control may be reduced.

Mixing Instructions

Fill the sprayer tank ½ full of water and start sprayer tank agitation. Add the required amount of Simplicity. Add the correct amount of Starane II Herbicide and continue to agitate.

Preharvest/Grazing Intervals (Starane II Herbicide and Simplicity)

- Livestock may be grazed on treated crops 7 days following application.
- Do not harvest the treated crop within 60 days after application.

Rates of Application

I. Low Rate: To control or suppress the following weeds apply a uniform spray containing Starane II Herbicide at 0.21 L/ha tank mixed with Simplicity at 375 mL/ha.

Weeds Controlled

cleavers (1-6 whorls)

wild oats (up to 4 leaf, 2 tiller) ♦

♦Under low wild oat populations (<75 plants/m²)

II. High Rate: To control the above weeds plus the additional weeds listed below, apply a uniform spray containing Starane II Herbicide at 0.31 L/ha with Simplicity at 500 mL/ha.

Additional Weeds Controlled

| | |
|---------------------------------|------------------------------------|
| barnyard grass (1-5 leaf) | kochia* (2-8 leaf) |
| chickweed, common (up to 10 cm) | red root pigweed (1-8 leaf) |
| cleavers (1-8 whorls) | smartweed (lady's thumb, 1-5 leaf) |
| foxtail, yellow (1-5 leaf) | volunteer canola** (1-6 leaf) |
| hemp nettle (1-8 leaf) | volunteer flax (1-12 cm) |
| Japanese brome (1-6 leaf) | |

*Including biotypes resistant to Group 2 herbicides that inhibit the ALS enzyme.

** will not control volunteer Imidazolinone-tolerant canola

Additional Weeds Suppressed

| | |
|------------------------------------|---------------------------|
| brome, downy (2-6 leaf, 2 tillers) | storks-bill (1-8 leaf) |
| foxtail, green (1-5 leaf) | wild buckwheat (1-4 leaf) |

TANK-MIX COMBINATIONS –STARANE II HERBICIDE + SIMPLICITY + OTHER HERBICIDES (Do not apply to oats)

When tank mixing Starane II Herbicide + Simplicity at either the low or high rates, the addition of other herbicides will provide control of additional weeds. For products which may be tank mixed with Starane II Herbicide + Simplicity and the additional weeds controlled see following table.

Tank-Mix Combinations with Starane II Herbicide (0.21 L/ha) + Simplicity Herbicide (375 mL/ha) or Starane II Herbicide (0.31 L/ha) + Simplicity herbicide (500 mL/ha)

| Herbicide Tank-Mix Partner | Crops Registered | Rate of Tank-Mix Partner | Additional Weeds Controlled |
|----------------------------|---|--------------------------|--|
| 2,4-D ester* | spring wheat, durum wheat, winter wheat | Up to 420 g ae/ha | Susceptible weeds such as: mustard (except dog and green tansy), flixweed, bluebur, burdock, cocklebur, stinkweed, goatsbeard, prickly lettuce, lamb's-quarters, plantain, ragweeds, Russian thistle, shepherd's purse, annual sunflower, sweet clover, volunteer canola, wild buckwheat, wild radish *Consult 2,4-D ester labels for weed stages and rates |
| MCPA ester* | spring wheat, durum wheat, winter wheat | Up to 560 g ae/ha | Burdock (before the 4-leaf stage), cocklebur, flixweed, plantain, lamb's-quarters, mustards (except dog and tansy), prickly lettuce, ragweeds, Russian pigweed, shepherd's-purse, stinkweed, vetch, wild buckwheat, wild radish, wild (annual) sunflower. *Consult MCPA ester labels for weed stages and rates |

RANGELAND, PERMANENT PASTURE, INDUSTRIAL AND OTHER NON-CROP AREAS OF CANADA**Starane II Herbicide Alone**

Apply Starane II Herbicide for post emergent control of kochia, including ALS resistant biotypes in non-cropland areas including roadsides, rights of way, permanent perennial grass cover areas, and industrial vegetation management areas. See Table for weed controlled and application rates.

Kochia Control by Starane II Herbicide Alone

| Weeds | Starane II Herbicide Rate (L/ha) |
|--------------------------------|----------------------------------|
| Kochia* (2-8 leaf) Control | 0.84 |
| Kochia* (2-8 leaf) Suppression | 0.42 |

*Including biotypes resistant to Group 2 herbicides that inhibit the ALS enzyme.

NOTE: Starane II Herbicide activity is influenced by weather conditions. Optimum activity requires active weed growth. The temperature range for optimum activity is 12°C to 24°C. Reduced activity will occur when temperatures are below 8°C or above 27°C. Frost before application (3 days) or shortly after (3 days) may reduce weed control. Weed control may be reduced during stress conditions, e.g. drought, heat or cold stress, or if weeds have initiated flowering, or if heavy infestations exist.

APPLICATION DIRECTIONS**Application Timing**

Apply when weeds are actively growing. Best results are obtained from applications made to seedling weeds. Only weeds emerged at the time of treatment will be controlled. Extreme growing conditions such as drought or near freezing temperature prior to, at and following time of application may reduce weed control. Foliage that is wet at the time of application may decrease control. Starane II Herbicide applications are rainfast within 1 hour after application.

Individual Plant or Spot/Strip Applications

Use the following mixing chart to mix Starane II Herbicide with enough water to make the correct amount of spray solution. Use equipment designed for individual plant or spot/strip application, such as a backpack sprayer or hose and handgun, to spray weeds. Thoroughly and uniformly wet the foliage of all target plants but not to the point of runoff. Direct spray away from aquatic habitats and non-target terrestrial plants.

Mixing Chart

| Starane II Herbicide | Spray Solution (water and herbicide) |
|----------------------|--------------------------------------|
| 4.2 – 8.4 mL | 10 L |
| 42 – 84 mL | 100 L |
| 420 – 840 mL | 1000 L |

Use the higher rate in the rate range for kochia control or when growing conditions are less than favourable. The maximum annual application rate is 840 mL/ha.

Restrictions

- **Grazing intervals:** There are no grazing restrictions for livestock following application.
- **Legumes:** Do not apply if the injury to legumes cannot be tolerated. Legumes may be less sensitive to herbicide injury after plant growth is mature and seed has set.
- Do not harvest grass for hay from treated areas within 7 days of application.

Tank Mix Combination – Starane II Herbicide + Milestone™ Herbicide

Starane II Herbicide can be tank mixed with Milestone Herbicide to broaden the spectrum of weeds and invasive plants controlled. Use 0.42 – 0.84 L/ha Starane II Herbicide with 0.25 – 0.5 L/ha Milestone Herbicide. For mixing order see mixing instructions.

Mixing Instructions

1. Begin to fill sprayer tank with clean water, and engage agitator. **Agitation must be continued throughout the entire mixing and spraying procedure.**

2. When the sprayer is three quarters full of water, add Milestone Herbicide. If more than one container of Milestone Herbicide is to be used, add all Milestone Herbicide containers prior to adding any other tank-mix partners.
3. Add Starane II Herbicide next. Continue to fill tank to desired level with water.

SPRAY BUFFER ZONES

A spray buffer zone is NOT required for:

- uses with hand-held equipment permitted on this label.

For application to rights-of-way, spray buffer zones for protection of sensitive terrestrial habitats are not required; however, the best available application strategies that minimize off-site drift, including meteorological conditions (for example, wind direction, low wind speed) and spray equipment (for example, coarse droplet sizes, minimizing height above canopy), should be used. However, applicators must observe the specified spray buffer zones for protection of sensitive aquatic habitats.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands) and sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

| Method of application | Crop | | Spray Buffer Zones (metres) Required for the Protection of: | | | | Terrestrial habitat |
|-----------------------|---|-------------|---|------------------|-------------------------------------|------------------|---------------------|
| | | | Freshwater Habitat of Depths: | | Estuarine/marine Habitat of Depths: | | |
| | | | Less than 1 m | Greater than 1 m | Less than 1 m | Greater than 1 m | |
| Field sprayer | Spring wheat, durum wheat, winter wheat, spring barley and oats | | 1 | 0 | 1 | 1 | 3 |
| | Rangeland and permanent pasture land, industrial and other non-crop areas | | 1 | 1 | 1 | 1 | 5* |
| Aerial | Spring wheat, durum wheat, winter wheat, spring barley and oats | Fixed Wing | 5 | 0 | 1 | 1 | 95 |
| | | Rotary Wing | 3 | 0 | 1 | 1 | 80 |

* Buffer zones for the protection of terrestrial habitats are not required for use on rights-of-way including railroad ballast, rail and hydro rights-of-way, utility easements, roads, and training grounds and firing ranges on military bases.

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticides portion of the Canada.ca website.

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, Starane II Herbicide is a Group 4 herbicide. Any weed population may contain or develop plants naturally resistant to Starane II Herbicide and other Group 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of Starane II Herbicide or other Group 4 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact your local Corteva Agriscience Canada Company representative or the Corteva Agriscience Canada Company hotline at 1-800-667-3852.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

The user assumes the risk to persons or property that arises from any such use of this product.

TMTrademark of Corteva Agriscience and its affiliated companies.

All other products listed are registered trademarks of their respective companies.