

# Distinct®

Herbicide

**Selective post-emergent weed control for field corn. Complements glyphosate for superior chemfallow and post-harvest control.**

- Controls annual broadleaf weeds in post-emergent corn
- Multiple modes of action with glyphosate to control resistant biotypes in chemfallow and post-harvest
- Help keep fields cleaner to set them up for success the next season
- Excellent follow-crop flexibility that includes pulses and canola

## Active ingredients

Dicamba – Group 4  
Diflufenzopyr – Group 19

## Formulation

Water dispersible granular

## One case contains

2 x 2.3 kg jugs

## Storage

Does not require heated storage.

### Weed control in spring, following previous September application



Source: BASF Small Plot Trials, 2013

### Glyphosate-resistant kochia 61 days after herbicide application



Source: BASF Small Plot Trials, 2013

## Application or crop

Field corn  
Chemfallow  
Post-harvest

## Staging

2 to 6 leaf  
July to August  
prior to first significant frost

## Weeds controlled and staging

### In field corn.

Apply to actively growing weeds (except where indicated)

Biennial wormwood (2 to 8 leaf)	Lamb's quarters
Canada thistle <sup>1</sup>	Perennial sow thistle <sup>2</sup> (2 to 10 leaf)
Common cocklebur (cotyledon to 6 leaf)	Redroot pigweed
Common ragweed	Tall waterhemp
Giant ragweed <sup>2</sup> (2 to 8 leaf)	Velvetleaf
Kochia (up to 15 cm height)	Volunteer adzuki beans (1 to 3 trifoliolate)
Lady's thumb	Volunteer canola (cotyledon to 4 leaf)
	Wild buckwheat

### In chemfallow and post-harvest.

Apply to actively growing weeds (except where indicated)

### Distinct at 58 g/ac tank mixed with glyphosate will control:

Dandelion <sup>1</sup>	Redroot pigweed
Kochia	Round-leaved mallow
Lamb's quarters	Spiny annual sow thistle
Narrow-leaved hawk's beard	Wild buckwheat

### Distinct at 115 g/ac tank mixed with glyphosate will control:

Biennial wormwood (2 to 8 leaf)	Lamb's quarters
Canada thistle <sup>1</sup>	Perennial sow thistle <sup>2</sup> (2 to 10 leaf)
Common cocklebur (cotyledon to 6 leaf)	Redroot pigweed
Common ragweed	Tall waterhemp
Dandelion <sup>1</sup>	Velvetleaf
Kochia <sup>3</sup> (up to 15 cm height)	Volunteer canola (cotyledon to 4 leaf)
Lady's thumb	Wild buckwheat

<sup>1</sup> Top growth.

<sup>2</sup> Suppression only.

<sup>3</sup> Includes glyphosate-resistant biotypes at 115 g/ac (285 g/ha) application rate.

**BASF**

We create chemistry

### Application rates

**One case of Distinct herbicide will treat 40 to 80 acres, depending on rate.**

Field corn	Distinct herbicide	115 g/ac (285 g/ha)
	Non-ionic surfactant <sup>4</sup>	0.25% v/v
	28% UAN <sup>4</sup>	1.25% v/v
Chemfallow, post-harvest	Distinct herbicide	58 to 115 g/ac (143 to 285 g/ha)
	Glyphosate <sup>4</sup> (360 g ae/L)	0.51 to 1 L/ac (1.25 to 2.5 L/ha)
	Merge adjuvant <sup>4</sup>	200 ml/ac (500 ml/ha)

### Water volume

Ground application only 20 to 40 L/ac (5 to 10 gal/ac)

<sup>4</sup> Non-ionic surfactant, 28% UAN, glyphosate and Merge adjuvant are not included in the case.

### Mixing order

1. Fill clean spray tank 1/2 full of clean water and start agitation.
2. Add the correct amount of Distinct herbicide and continue to agitate until product is completely dissolved and fully dispersed.
3. For chemfallow or post-harvest, add the correct amount of glyphosate while continuing agitation.
4. For chemfallow or post-harvest, add the recommended amount of Merge adjuvant or a 0.25% v/v of non-ionic surfactant, followed by 1.25% v/v 28% UAN. For field corn, add 0.25% v/v of non-ionic surfactant, followed by 1.25% v/v 28% UAN.
5. Continue agitation while adding the remaining amount of water.

### Application tips

**Rainfastness** – 4 hours.

For denser weeds and thick canopies, use the higher water volume.

### Grazing

Do not cut or graze corn for 75 days after application. See label for additional restrictions.

### Follow crops

#### If Distinct is applied prior to September 1

Wheat, barley, oats, canary seed, corn, canola, lentils, soybeans, chickpeas, flax, field peas and sunflowers.

#### If Distinct is applied<sup>5</sup> prior to October 1

Wheat, barley, oats, canary seed, corn, canola, lentils, field peas and soybeans.

#### If Distinct is applied<sup>5</sup> prior to October 15

Wheat, barley, oats, canary seed and corn.

<sup>5</sup> Distinct applied at 58 g/ac (143 g/ha). If higher rate is used, rotate to cereal or corn crops only.

### Tank mixes

#### Herbicide for chemfallow and post-harvest application: Glyphosate

None on label for field corn.

Contact your local BASF **AgSolutions**® Grower or Retail Representative or call **AgSolutions** Customer Care at 1-877-371-BASF (2273) for additional information on supported tank mixes.

**For more information:** Call **AgSolutions** Customer Care at 1-877-371-BASF (2273) or visit [agsolutions.ca](https://www.agsolutions.ca).

### Always read and follow label directions.

**AgSolutions**, **DISTINCT** and **MERGE** are registered trademarks of BASF; all used under license by BASF Canada Inc. © 2024 BASF Canada Inc.