

Voraxor®

Powered by **Tirexor®** Herbicide

Active ingredients Saflufenacil and Trifludimoxazin – Group 14

One case contains 1 x 1.56 L jug of Voraxor® herbicide
2 x 8.1 L jugs of Merge® adjuvant

Crops

Barley
Chickpeas
Faba beans
Field corn
Lentils
Peas (dried field)
Soybeans
Wheat (including durum, spring and winter)

Staging

pre-seed and pre-emergence

Chemfallow

Apply to actively growing weeds
less than 15 cm in height

Weeds controlled

Broadleaves¹

Canada fleabane

Cleavers²

Kochia²

Lamb's quarters²

Narrow-leaved hawk's beard

Redroot pigweed²

Round-leaved mallow

Shepherd's-purse³

Stinkweed²

Volunteer canola^{2,4}

Wild buckwheat²

Wild mustard²

Application rates⁴

One case of Voraxor herbicide will treat 27 to 80 acres, depending on rate. One case will treat 80 acres for burndown and 30 to 40 acres for residual activity.

Lentils

19.5 ml/ac (48 ml/ha)⁵

Field corn and soybeans

19.5 – 40.5 ml/ac (48 – 100 ml/ha)⁶

Wheat (including durum, spring and winter), peas, chickpeas, faba beans and barley

19.5 – 58 ml/ac (48 – 144 ml/ha)

Chemfallow

19.5 – 29 ml/ac (48 – 72 ml/ha)

All applications

Merge adjuvant

200 to 400 ml/ac (0.5 to 1 L/ha)

Water volume

Ground application

50 to 100 L/ha (13 gal/ac)

 **BASF**

We create chemistry

Application tips

- Restricted entry interval – 12 hours.
- Do not apply during periods of dead calm or when winds are gusty.
- Should the product freeze, thaw at room temperature and agitate well before use.

Mixing order

1. Thoroughly clean the sprayer prior to use by flushing the system with water containing detergent. Refer to previously applied product labels for specific cleaning instructions.
2. Fill the clean spray tank $\frac{1}{2}$ full with clean water. Start the agitation system. Agitation should be running during the entire mixing procedure.
3. Add the correct amount of Voraxor and continue to agitate until thoroughly mixed.
4. Prior to adding glyphosate, add the correct amount of herbicide tank mix partner if required.
5. Add in glyphosate.
6. Add the correct amount of Merge adjuvant.
7. Continue agitation while filling the remainder of the tank with water necessary to fill the spray tank.
8. Maintain continuous and constant agitation throughout application until spraying is complete. After any break in spraying, agitate thoroughly before spraying again.
9. Thoroughly clean the sprayer after use by flushing the system with clean water and detergent.

Tank mixes

Herbicide for all crops: glyphosate

Herbicide for lentils, field corn, soybeans, peas (dried field), chickpeas and faba beans: Zidua[®] SC herbicide

For more information about Voraxor herbicide, please call your BASF Sales Representative, visit agsolutions.ca/Voraxor or call **AgSolutions[®]** Customer Care at 1-877-371-BASF (2273).

¹ Voraxor applied at 19.5 – 29 ml/ac (48 – 72 ml/ha) provides rapid burndown control of all weeds listed below.

² Apply Voraxor at a rate of 41.5 – 57 ml/ac (100 – 140 ml/ha) for suppression of secondary flushes.

³ Suppression only.

⁴ All types including Roundup Ready[®].

⁵ Rate restrictions apply. Do not use rate higher than 19.5 ml/ac (48 ml/ha) or injury could result.

⁶ Rate restrictions apply. Do not use higher than 40.5 ml/ac (100 ml/ha) or injury could result.

Always read and follow label directions.

AgSolutions, **MERGE**, **TIREXOR**, **VORAXOR** and **ZIDUA** are registered trademarks of BASF; all used under license by BASF Canada Inc. © 2023 BASF Canada Inc.

Roundup Ready[®] is a registered trademark of Bayer Group. Used under license.

© 2023 Bayer Group. All rights reserved.

Voraxor[®]

Powered by **Tirexor[®]** Herbicide