

# Voraxor®

Powered by **Tirexor®** Herbicide

SUPPORTED BY THE

**ADVANCED WEED CONTROL**

PROGRAM

## The new standard in pre-seed burndown control.

- Improved burndown with increased activity, compared to Heat® LQ herbicide, on most broadleaf weeds
- Enhanced residual consistency and weed spectrum, including cleavers, kochia and wild mustard
- Contains the new unique active ingredient Tirexor®

### Active ingredient

Saflufenacil – Group 14  
Trifludimoxazin – Group 14

### Formulation

Suspension concentrate


### One case contains

1 x 1.56 L jug Voraxor® herbicide  
2 x 8.1 L Merge® adjuvant

### Storage

Requires heated storage.

### Voraxor vs competitors, 14 to 21 days after treatment (DAT)



Glyphosate was not included in this trial.  
Source: BASF Small Plot Trials, Saskatoon, SK, 2020

### Crops

Barley  
Chickpeas  
Faba beans  
Field corn  
Lentils  
Peas (dried field)  
Soybeans  
Wheat (incl. durum, spring and winter)  
Chemfallow

### Staging

pre-seed and pre-emergence

apply to actively growing weeds less than 15 cm in height

### Weeds controlled

#### Broadleaves<sup>1</sup>

Canada fleabane  
Cleavers<sup>2</sup>  
Kochia<sup>2</sup>  
Lamb's quarters<sup>2</sup>  
Narrow-leaved hawk's beard  
Redroot pigweed<sup>2</sup>  
Round-leaved mallow  
Shepherd's-purse<sup>3</sup>  
Stinkweed<sup>2</sup>  
Volunteer canola<sup>2,4</sup>  
Wild buckwheat<sup>2</sup>  
Wild mustard<sup>2</sup>

### Staging

8 leaf (except where indicated)

(4 whorl stage)  
(15 cm)

(8 cm)

### Application rates<sup>4</sup>

**One case of Voraxor herbicide will treat 30 to 80 acres, depending on rate. One case will treat 80 acres for burndown and 30 to 40 acres for residual control.**

Lentils	19.5 ml/ac (48 ml/ha) <sup>5</sup>
Field corn and soybeans	19.5 to 40.5 ml/ac (48 to 100 ml/ha) <sup>6</sup>
Faba beans, chickpeas, wheat (incl. durum, spring and winter), peas and barley	19.5 to 58 ml/ac (48 to 144 ml/ha)
Chemfallow	19.5 to 29 ml/ac (48 to 72 ml/ha)

### All applications

Merge adjuvant 200 to 400 ml/ac (0.5 to 1 L/ha)

### Water volume

Ground application only 20 to 40 L/ac (5 to 10 gal/ac)

<sup>1</sup> Voraxor applied at 19.5 to 29 ml/ac (48 to 72 ml/ha) provides rapid burndown control of all weeds listed below.

<sup>2</sup> Apply Voraxor at a rate of 41.5 to 57 ml/ac (100 to 140 ml/ha) for suppression of secondary flushes.

<sup>3</sup> Suppression only.

<sup>4</sup> All types including Roundup Ready®.

<sup>5</sup> Rate restrictions apply. Do not use rate higher than 19.5 ml/ac (48 ml/ha) or injury could result.

<sup>6</sup> Rate restrictions apply. Do not use higher than 40.5 ml/ac (100 ml/ha) or injury could result.


**BASF**

We create chemistry

# Voraxor®

Powered by **Tirexor®** Herbicide


**Annual broadleaf weed control burndown, 28 to 36 DAT**



Weeds controlled: redroot pigweed, volunteer canola, lamb's quarters, wild mustard, cleavers, kochia and wild buckwheat

Source: BASF Small Plot Trials, 2020, n=22

**New active with improved burndown over Heat LQ**



Glyphosate was not included in this trial.

Source: BASF Small Plot Trials, 2020-2021, n=25 to 28 DAT

## For more information:

Call **AgSolutions** Customer Care at 1-877-371-BASF (2273)

Visit **agsolutions.ca**

## Mixing order

1. Thoroughly clean the sprayer prior to use by flushing the system with water containing detergent. Refer to previously applied product labels for specific cleaning instructions.
2. Fill the clean spray tank 1/2 full with clean water. Start the agitation system. Agitation should be running during the entire mixing procedure.
3. Add the correct amount of Voraxor and continue to agitate until thoroughly mixed.
4. Prior to adding glyphosate, add the correct amount of herbicide tank-mix partner if required.
5. Add in glyphosate.
6. Add the correct amount of Merge adjuvant.
7. Continue agitation while filling the remainder of the tank with water necessary to fill the spray tank.
8. Maintain continuous and constant agitation throughout application until spraying is complete. After any break in spraying, agitate thoroughly before spraying again.
9. Thoroughly clean the sprayer after use by flushing the system with clean water and detergent.

## Application tips

**Restricted entry interval** –12 hours.

Avoid application when heavy rain is forecast.

Do not apply during periods of dead calm or when winds are gusty.

## Pre-harvest interval

There is no required pre-harvest interval between a pre-seed or pre-emergent application and harvest.

## Follow crops

**3 months after application**<sup>7</sup>

Winter wheat

**Plant back crops**<sup>8</sup>

Barley, dry field peas, lentils, field corn, soybean, wheat (spring, winter, durum)

**Following season**<sup>7</sup>

Barley, canola, dry common beans, dry field peas, flax, field corn, lentils, mustard, soybean, wheat (spring, durum)

<sup>7</sup> Following a spring application of Voraxor.

<sup>8</sup> To be planted in the same season in case of crop failure. Rate restrictions apply. Lentils, field corn and soybeans can only be grown as plant back crops provided that a maximum product rate of 19.5 ml/ac (48 ml/ha) for lentils and 40.5 ml/ac (100 ml/ha) for field corn and soybeans was applied in the previous crop. A second application of Voraxor cannot be made in the rescue crop. Crops can also be planted in the next season following chemfallow treatment applied after August 1.

## Tank mixes

**Herbicide for all crops:** Glyphosate

**Herbicide for lentils, field corn, soybeans, peas (dried field), chickpeas and faba beans:**

Zidua® SC herbicide

Contact your local BASF **AgSolutions®** Grower or Retail Representative or call **AgSolutions** Customer Care at 1-877-371-BASF (2273) for additional information on supported tank mixes.

## Always read and follow label directions.

**AgSolutions**, HEAT, MERGE, TIREXOR, VORAXOR and ZIDUA are registered trademarks of BASF; all used under license by BASF Canada Inc. © 2023 BASF Canada Inc.

Aim, Authority, and Focus are registered trademarks of FMC Corporation or an affiliate. Intruvix is a trademark of FMC Corporation or an affiliate.

BlackHawk and GoldWing are registered trademarks of Nufarm Agriculture Inc.

Roundup Ready® is a registered trademark of Bayer Group. Used under license. © 2024 Bayer Group. All rights reserved.